


Mortality down with spinal anesthesia for hip Fx surgery

April 20 2018



(HealthDay)—For patients undergoing hip fracture fixation, general

anesthesia (GA) is associated with increased 90-day mortality compared with spinal anesthesia (SA), according to a study presented at the 2018 World Congress on Regional Anesthesia and Pain Medicine, held from April 19 to 21 in New York City.

Sarah Tierney, M.D., from the University of Toronto, and colleagues conducted a retrospective study to examine the correlation of anesthetic technique and mortality in patients undergoing hip fracture fixation over a 13-year period. A total of 2,591 patients who underwent hip fracture fixation were identified; 883 of these in the SA group were matched to patients in the GA group. Clinical, laboratory, and outcome data were obtained, and [mortality data](#) were collected from the hospital discharge database.

The researchers found that patients receiving GA had increased 90-day mortality (8.6 versus 6.3 percent; P pulmonary embolism and major blood loss (2.3 versus 0.5 percent and 7.7 versus 4.8 percent, respectively; both P

"This observational propensity-score, matched-pairs, cohort study suggests a strong association between spinal anesthesia and lower 90-day mortality, as well as reduced rates of pulmonary embolism [and] major blood loss and a shorter hospital length of stay," the authors write.

More information: [More Information](#)

Copyright © 2018 [HealthDay](#). All rights reserved.

Citation: Mortality down with spinal anesthesia for hip Fx surgery (2018, April 20) retrieved 27 April 2024 from

<https://medicalxpress.com/news/2018-04-mortality-spinal-anesthesia-hip-fx.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.