

Nemours study highlights psychological and social barriers to treating childhood obesity

April 5 2018

Children whose families have elevated psychological and social risks, including child behavior problems, parent mental health issues, and family financial difficulties, were more likely to drop out of weight management treatment and less likely to have an improvement in weight status, according to a study published online today by the *Journal of Pediatrics*. The study, a collaboration between researchers at the Nemours Center for Healthcare Delivery Science and Nemours Division of Weight Management, supports the need for psychosocial screening early in the treatment of childhood obesity.

"Previous studies have found that the majority of children who receive interdisciplinary treatment in [weight management](#) clinics are successful at achieving a healthier [weight](#). However, most weight management clinics report significant drop-out rates, limiting the number of patients who benefit from treatment," said Thao-Ly Tam Phan, MD, MPH, a pediatrician in the Nemours Weight Management Clinic, and lead author of the study. "Given the significant impact that [psychosocial risk factors](#) seem to have on outcomes in weight management treatment, identifying and addressing parent mental health concerns, child behavior concerns, and family social resource needs upfront may help improve outcomes in children with obesity."

The prospective study, funded by the Academic Pediatric Association's Research in Academic Pediatrics Initiative on Diversity (RAPID) Award received by Dr. Phan, enrolled 100 families of children between the ages of 4 and 12 during their first visit to the weight management clinic. The

majority of children were non-Hispanic Black (36 percent) or White (43 percent) and had severe obesity (55 percent). Researchers found that 59 percent of families had moderate to high psychosocial [risk](#) scores and 41 percent had low scores on the Psychosocial Assessment Tool (PAT), a validated screener of family psychosocial risk. Compared to data on all patients who have completed the PAT in the United States and internationally, families in this study were two times more likely to have a moderate-to-high risk score.

Each child's progress was tracked over six months, and the study found that families with moderate-to-high risk scores on the PAT were 3.1 times more likely to stop attending the clinic. Additionally, children of those families were 2.6 times more likely to have an increase in body mass index (BMI) z-score and 3.2 times more likely to not have a meaningful change in their BMI z-score. Notably, 14 patients receiving psychological services as a component of their weight [management](#) treatment experienced lower attrition rates than those who did not (27 percent vs. 57 percent).

"These findings together highlight how important [family](#) psychosocial factors are to a child's health behaviors and weight," said Anne E. Kazak, Ph.D., ABPP, the co-director of the Center for Healthcare Delivery Science at Nemours Children's Health System and creator of the PAT.

The authors suggest that future studies evaluate the long-term impact of psychosocial risk and of providing psychosocial services to families of children with obesity. Through a National Institute of Child Health and Human Development-sponsored K23 Mentored Patient-Oriented Research Career Development Award, Drs. Phan and Kazak are taking this next step by developing and testing an innovative intervention to provide parents with guidance on managing their child's behaviors and to engage families of diverse backgrounds in obesity [treatment](#).

More information: Thao-Ly T. Phan, MD, MPH, et al. "Impact of Psychosocial Risk on Outcomes among Families Seeking Treatment for Obesity." *Journal of Pediatrics* (2018): April 5, 2018. Web.

Provided by Nemours Children's Health System

Citation: Nemours study highlights psychological and social barriers to treating childhood obesity (2018, April 5) retrieved 2 May 2024 from <https://medicalxpress.com/news/2018-04-nemours-highlights-psychological-social-barriers.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.