


Pregnant women are at increased risk of domestic violence in all cultural groups

April 26 2018, by Hannah Dahlen And Virginia Schmied



Credit: AI-generated image ([disclaimer](#))

Domestic violence occurs across all age groups and life stages. Rather than reducing during pregnancy, expecting a child is a [key risk factor for domestic violence beginning or escalating](#).

Our research, [published today in the journal BMJ Open](#), found that 4.3%

of pregnant women due to give birth in Western Sydney disclosed [domestic violence](#) when asked about it by a midwife at her first hospital visit. The study examined more than 33,000 ethnically diverse women who gave birth between 2006 and 2016, and found that these disclosures spanned all cultural groups.

Domestic [violence](#) in pregnancy not only causes distress and trauma for the mother and baby, it increases the risk of the baby having a [low birth weight](#) (very small baby) or being born prematurely (before 37 weeks), which is linked to jaundice, anaemia and respiratory distress in infancy, and diabetes and heart disease later in life.

Abuse and trauma

Depending on the state or territory, women may receive a ["psychosocial" assessment](#) from midwives when they first book into a public hospital during pregnancy. This screens for depression, anxiety, childhood [abuse](#), domestic violence, support and stress.

Using these assessments, we found that 4.3% of women disclosed domestic violence overall, but rates were higher among women having a subsequent baby, compared with first-time mothers.

We're unsure if this is because violence has escalated for these women with subsequent pregnancies; if they trust health providers more to disclose the violence; or if they seek help because they're becoming more aware of the impact of domestic violence on their children.

We found a strong link between the disclosure of childhood abuse and the disclosure of domestic violence. Nearly 24% of women disclosing domestic violence had also disclosed childhood abuse.

This doesn't mean that one causes the other, but women who experience

[childhood abuse](#) are more vulnerable to re-victimisation (being abused again). They may feel like they're not worthy and gravitate towards men who abuse them.

Women who disclosed domestic violence in our study were more likely to have a history of anxiety or depression (34%) and have thoughts of harming themselves.

This is concerning, as maternal suicide during pregnancy or following the birth appears to be rising and has now become one of the main causes of maternal death in Australia.

We also found rising rates of [pregnant women](#) disclosing domestic violence and being admitted to hospital for bleeding and signs that labour may be starting early (before 37 weeks). When this happens, women are admitted to hospital to try to stop the labour, or to find the source of the bleeding. Sometimes stress can [contribute](#) to preterm birth and bleeding in pregnancy.

Ethnic backgrounds

We found that domestic violence occurred across all cultural groups, but reported rates were highest among women from New Zealand and Sudan.

Previous research has shown high rates of domestic violence among Maori women in some parts of [New Zealand](#).

There is also evidence of high rates of domestic violence among [Sudanese](#) women prior to migration.

We found that women born in India and China reported very low rates of domestic violence. This may reflect a cultural tendency not to discuss

what is considered private family business with outsiders.

It's important that health professionals know how to ask about domestic violence in a culturally appropriate way so women feel comfortable disclosing abuse and can access appropriate support.

What needs to be done?

Midwives need to consider cultural norms and acceptability when asking migrant women questions about domestic violence, and this must always be done in a way that keeps the woman safe. Partners should not be present when the questions are asked – and this may be done at another time in the pregnancy if necessary.

Where English is not the first language, interpreters should be used. But this can also present challenges if the interpreter comes from the same community and is known to the woman.

When women have continuity of midwifery care and get to know a midwife well throughout the [pregnancy](#), it is easier for midwives to gain [women's](#) trust and to notice when things change. This style of care should be rolled out more widely in Australian public hospitals.

This article was originally published on [The Conversation](#). Read the [original article](#).

Provided by The Conversation

Citation: Pregnant women are at increased risk of domestic violence in all cultural groups (2018, April 26) retrieved 23 April 2024 from <https://medicalxpress.com/news/2018-04-pregnant-women-domestic-violence-cultural.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.