


Scientists induce sensations of touch and movement in the arm of a paralyzed man

April 10 2018



Primary Somatosensory
cortex activation

fMRI is used to highlight select implant sites in the somatosensory cortex. Electrodes implanted in this region were able to stimulate neurons that produced physical sensations, like a squeeze or tap, in the arm of a paralyzed man. Credit: Andersen lab

For the first time, scientists at Caltech have induced natural sensations in the arm of a paralyzed man by stimulating a certain region of the brain with a tiny array of electrodes. The patient has a high-level spinal cord lesion and, besides not being able to move his limbs, also cannot feel them. The work could one day allow paralyzed people using prosthetic limbs to feel physical feedback from sensors placed on these devices.

The research was done in the laboratory of Richard Andersen, James G. Boswell Professor of Neuroscience, T&C Chen Brain-Machine Interface Center Leadership Chair, and director of the T&C Chen Brain-Machine Interface Center. A paper describing the work appears in the April 10 issue of the journal *eLife*.

The [somatosensory cortex](#) is a strip of brain that governs bodily sensations, both proprioceptive sensations (sensations of movement or the body's position in space) and cutaneous sensations (those of pressure, vibration, touch, and the like). Previous to the new work, neural implants targeting similar brain areas predominantly produced sensations such as tingling or buzzing in the hand. The Andersen lab's implant is able to produce much more natural [sensation](#) via intracortical stimulation, akin to sensations experienced by the patient prior to his injury.

The patient had become paralyzed from the shoulders down three years ago after a spinal cord injury. Two arrays of tiny electrodes were surgically inserted into his somatosensory cortex. Using the arrays, the researchers stimulated neurons in the region with very small pulses of electricity. The participant reported feeling different natural sensations—such as squeezing, tapping, a sense of upward motion, and several others—that would vary in type, intensity, and location depending on the frequency, amplitude, and location of stimulation from the arrays. It is the first time such natural sensations have been induced by intracortical neural stimulation.

"It was quite interesting," the study participant says of the sensations. "It was a lot of pinching, squeezing, movements, things like that. Hopefully it helps somebody in the future."

Though different types of stimulation did indeed induce varying sensations, the neural codes governing specific physical sensations are still unclear. In future work, the researchers hope to determine the precise ways to place the electrodes and stimulate somatosensory brain areas in order to induce specific feelings and create a kind of dictionary of stimulations and their corresponding sensations.

The next major step, according to Andersen, is to integrate the technology with existing neural prosthetics. In 2015, Andersen's laboratory developed brain-machine interfaces (BMIs) to connect a prosthetic robotic arm to electrodes implanted in the region of the brain that governs intentions. In this way, a paralyzed man was able to utilize the prosthetic arm to reach out, grasp a cup, and bring it to his mouth to take a drink. Connecting the device with the somatosensory cortex would create bidirectional BMIs that would enable a paralyzed person to feel again, while using [prosthetic limbs](#).

"Currently the only feedback that is available for neural prosthetics is visual, meaning that participants can watch the brain-controlled operation of robotic limbs to make corrections," says Andersen. "However, once an object is grasped, it is essential to also have somatosensory information to dexterously manipulate the object. Stimulation-induced somatosensory sensations have the potential added advantage of producing a sense of embodiment; for example, a participant may feel over time that the robotic limb is a part of their body."

The study is titled "Proprioceptive and Cutaneous Sensations in Humans Elicited by Intracortical Microstimulation."

More information: Michelle Armenta Salas et al, Proprioceptive and cutaneous sensations in humans elicited by intracortical microstimulation, *eLife* (2018). [DOI: 10.7554/eLife.32904](https://doi.org/10.7554/eLife.32904)

Provided by California Institute of Technology

Citation: Scientists induce sensations of touch and movement in the arm of a paralyzed man (2018, April 10) retrieved 3 May 2024 from <https://medicalxpress.com/news/2018-04-scientists-sensations-movement-arm-paralyzed.html>

| |
|--|
| <p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p> |
|--|