

# Temperature affects insecticide efficacy against malaria vectors

April 16 2018

---



Malaria-transmitting *Anopheles* mosquitoes are becoming increasingly resistant to widely used insecticides. Credit: ISGlobal

Ambient temperature has a marked effect on the toxicity of the most commonly used insecticides for malaria control, according to a study led by ISGlobal. The results, published in the *Malaria Journal*, underline the need to evaluate the efficacy of these chemicals under real field conditions.

The appearance and spread of mosquitoes resistant to the insecticides

currently used for [malaria control](#) is a threat to malaria elimination efforts. Pyrethroids are the only insecticide class approved for treating bed nets and used widely for indoor spraying. In Africa, resistance to pyrethroids has been reported. It is therefore crucial to monitor insecticide susceptibility or resistance among the main species of malaria-transmitting Anopheles. Insecticide efficacy is not only determined by the active ingredient, but also by other factors, including [ambient temperature](#). However, susceptibility tests are normally performed in laboratories or insectaries where temperature conditions are optimal for the mosquito.

In this study, the researchers explored the effect of temperature on the standard [insecticide](#) resistance test using resistant or susceptible strains of two major [malaria](#) vectors, *An. arabiensis* and *An. funestus*. Toxicity of the pyrethroid deltamethrin and the carbamate bendiocarb was assessed at different temperatures (18, 25 or 30 degrees).

Results show that temperature impacts toxicity of both insecticides, but in a different way—bendiocarb lost efficacy at higher temperatures for both species whether they were resistant or susceptible. In contrast, higher temperatures decreased deltamethrin [toxicity](#) for susceptible *arabiensis*; the contrary was observed for resistant *An. arabiensis* and susceptible *An. funestus*. Piperonyl-butoxide (PBO), which inhibits pyrethroid-resistance mechanisms, completely restored deltamethrin susceptibility at all temperatures.

The authors conclude that caution must be exercised when drawing conclusions about a chemical's efficacy from laboratory assays performed at only one [temperature](#). Temperatures in the field, they point out, can vary considerably during a single day.

"Performing efficacy tests with local vectors and under real field conditions (which would reflect the appropriate season and relevant time

of day when chemicals are expected to act) would yield more accurate entomological intelligence for evidence-based decision-making," says ISGlobal researcher Krijn Paaijmans, who coordinated the study.

**More information:** Katey D. Glunt et al, The impact of temperature on insecticide toxicity against the malaria vectors *Anopheles arabiensis* and *Anopheles funestus*, *Malaria Journal* (2018). [DOI: 10.1186/s12936-018-2250-4](https://doi.org/10.1186/s12936-018-2250-4)

Provided by Barcelona Institute for Global Health

Citation: Temperature affects insecticide efficacy against malaria vectors (2018, April 16) retrieved 12 September 2024 from <https://medicalxpress.com/news/2018-04-temperature-affects-insecticide-efficacy-malaria.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.