

Under-fives should be priority for snail fever

April 17 2018

Infants in some of the world's poorest regions are vulnerable to a common worm parasite infection and their treatment should become a priority, according to a study.

Regular testing and treatment of pre-school children for [snail fever](#) - known as schistosomiasis or bilharzia - would reduce the spread of the disease, while promoting [childhood health](#) and development, experts have found.

The parasite, which is carried by freshwater snails, mainly infects people in sub-Saharan Africa. Once diagnosed, treatment with a cheap, readily available drug is usually effective.

Common symptoms include abdominal pain or blood in the urine. In children, it can cause stunted growth and learning difficulties. Infection acquired in childhood can cause bladder cancer and liver damage, or impact on reproductive health, in adults.

Until now, pre-school children have not been routinely tested for nor treated for the condition, as the health impact of childhood infection has not been well understood.

In the first long-term study of the disease in this age group, researchers at the University of Edinburgh and collaborators in Zimbabwe monitored 1500 children aged from six months to five years old over the course of a year.

Their research showed that children could contract an infection within their first year of life, and develop symptoms within three months. The study also highlighted for the first time that infected children are at risk of malnutrition and stunted growth. Treatment of the infection with the drug Praziquantel, was shown to reverse symptoms.

The study, funded by Wellcome and Thrasher Research Fund, was published in the *BMJ Global Health*.

Professor Francisca Mutapi of the University of Edinburgh's School of Biological Sciences, who led the study, said: "We now know there is no such thing as snail fever without symptoms, and [young children](#) can get infected by snail fever very early in their lives."

Derick Osakunor, of the University of Edinburgh's School of Biological Sciences, who took part in the project, said: "Our findings show that young children can develop clinical disease quickly, but existing diagnosis methods and treatments are effective in under-fives, which is welcome news for managing cases of [infection](#) in young [children](#)."

More information: Derick Nii Mensah Osakunor et al, Dynamics of paediatric urogenital schistosome infection, morbidity and treatment: a longitudinal study among preschool children in Zimbabwe, *BMJ Global Health* (2018). [DOI: 10.1136/bmjgh-2017-000661](https://doi.org/10.1136/bmjgh-2017-000661)

Provided by University of Edinburgh

Citation: Under-fives should be priority for snail fever (2018, April 17) retrieved 7 May 2024 from <https://medicalxpress.com/news/2018-04-under-fives-priority-snail-fever.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.