

Women's wellness: Understanding depression and the gender gap

April 17 2018, by From Mayo Clinic News Network, Mayo Clinic News Network

Women are nearly twice as likely as men to be diagnosed with depression. Depression can occur at any age.

Some mood changes and depressed feelings occur with normal hormonal changes. But hormonal changes alone don't cause [depression](#). Other biological factors, inherited traits, and personal life circumstances and experiences are associated with a higher risk of depression. Here's what contributes to depression in women.

PUBERTY

Hormone changes during puberty may increase some girls' risk of developing depression. However, temporary mood swings related to fluctuating hormones during puberty are normal—these changes alone don't cause depression.

Puberty is often associated with other experiences that can play a role in depression, such as:

- Emerging sexuality and identity issues—Conflicts with parents—Increasing pressure to achieve in school, sports or other areas of life

After puberty, depression rates are higher in females than in males. Because girls typically reach [puberty](#) before boys do, they're more likely

to develop depression at an earlier age than boys are. This depression gender gap lasts until after menopause.

PREMENSTRUAL PROBLEMS

For most females with premenstrual syndrome (PMS), symptoms such as abdominal bloating, breast tenderness, headache, anxiety, irritability and experiencing the blues are minor and short-lived. But a small number of females have severe and disabling symptoms that disrupt their studies, jobs, relationships or other areas of their lives. At that point, PMS may cross the line into [premenstrual dysphoric disorder](#) (PMDD) - a type of depression that generally requires treatment.

The exact interaction between depression and PMS remains unclear. It's possible that cyclical changes in estrogen, progesterone and other hormones can disrupt the function of brain chemicals such as serotonin that control mood. Inherited traits, life experiences and other factors appear to play a role.

PREGNANCY

Dramatic [hormonal changes](#) occur during pregnancy, and these can affect mood. Other issues also may increase the risk of developing depression during pregnancy or during attempts to become pregnant, such as:

- Lifestyle or work changes or other life stressors—Relationship problems—Previous episodes of depression, postpartum depression or PMDD—Lack of social support—Unintended or unwanted pregnancy—Miscarriage—Infertility—Stopping use of antidepressant medications

POSTPARTUM DEPRESSION

Many new mothers find themselves sad, angry and irritable, and experience crying spells soon after giving birth. These feelings—sometimes called the baby blues—are normal and generally subside within a week or two. But more-serious or long-lasting depressed feelings may indicate [postpartum depression](#), particularly if signs and symptoms include:

- Low self-esteem or feeling like you're a bad mom—Anxiety or feeling numb—Trouble sleeping, even when your baby is sleeping—Problems with daily functioning—Inability to care for your baby—Thoughts of harming your baby—Thoughts of suicide

Postpartum depression is a serious medical condition requiring prompt treatment. It occurs in about 10 to 15 percent of women. It's thought to be associated with:

- Major hormonal fluctuations that influence mood—The responsibility of caring for a newborn—Predisposition to [mood](#) and anxiety disorders—Birth complications—Breast-feeding problems

- Infant complications or special needs—Poor social support

©2018 Mayo Foundation for Medical Education and Research
Distributed by Tribune Content Agency, LLC.

Citation: Women's wellness: Understanding depression and the gender gap (2018, April 17) retrieved 11 May 2024 from <https://medicalxpress.com/news/2018-04-women-wellness-depression-gender-gap.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.