


# A new way to watch brain activity in action

May 7 2018, by Katherine Fenz

---



It's a neuroscientist's dream: being able to track the millions of interactions among brain cells in animals that move about freely, behaving as they would under natural circumstances. New technology developed at The Rockefeller University represents a big step toward realizing that goal.

The invention, reported today in *Nature Methods*, is expected to give

researchers a dynamic tool to study the [brain](#)'s role in various behaviors. Although it is designed for use on mice, information gleaned from it could someday shed light on neuronal activity in humans as well, says Alipasha Vaziri, who led the technology's development as head of the Laboratory of Neurotechnology and Biophysics. For example, it might allow us to better understand the neuronal basis of brain disorders such as autism and schizophrenia.

Vaziri says the tool provides an opening to an exciting range of discoveries. As an animal moves about its environment, for instance, some neurons direct spatial navigation while others receive sensory feedback from changes to the body's position or the visual system. "Until now, no one has been able to detect how these different neurons, which can be located at different depths within a volume of brain tissue, dynamically interact with each other in a freely moving rodent," says Vaziri, an associate professor at Rockefeller. Similarly, the tool can be used to record the interplay among neurons when two animals meet and interact socially.

## High-tech headgear

The technology consists of a tiny microscope attached to a mouse's head and outfitted with a specialized group of lenses called a microlens array. These lenses enable the microscope to capture images from multiple angles and depths on a sensor chip, producing a three-dimensional record of neurons blinking on and off as they communicate with each other through electrochemical impulses. (In the experiments, the mouse neurons are genetically modified to light up when they become activated.) A coaxial cable attached to the top of the microscope transmits the data for recording. The head-mounted gear weighs about four grams, about as much as a mouse can support, but Vaziri expects that planned modifications will make it significantly lighter.

Once the microlens array has captured sensor images from within a volume of [brain tissue](#), the next challenge is to process this raw data. Brain tissue is opaque, making it difficult to pinpoint the source of each neuronal light flash. Vaziri's team solved this problem, which is the result of so-called scattering, by developing a new computer algorithm. "The algorithm utilizes the statistical properties of neurons' distribution in space and in activity," Vaziri explains, "while extracting additional information from the scattered emission light. This enables their activity to be simultaneously and faithfully recorded within a volume despite of the highly scattering tissue properties."

The result is a clear image that shows individual [neurons](#) flashing in sequence.

## **Faster, more effective imaging**

Vaziri's lab has previously applied this [algorithm, known by the acronym SID](#), in studies in which the heads of the mice were secured in a fixed position. Their latest research is the first to demonstrate that these inventions can be used together with a tiny microscope called the Miniscope, developed by a collaborating team at the University of California Los Angeles, to measure neuronal activity volumetrically in unconstrained animals.

The technology, if widely adopted, could offer several advantages over two-photon microscopy, a broadly used neuroscience tool. For example, two-photon microscopy records neuronal activity within individual focal planes—thin, virtual "slices" of the sample—that then are combined to create a three-dimensional image. In contrast, Vaziri's method immediately captures data in three dimensions over an entire volume of tissue, making it faster and more effective.

Vaziri plans to continue developing tools to record [neuronal activity](#) in

even larger portions of the brain than is currently possible, and at higher speeds and resolution. "We hope this work will ultimately lead to a deeper understanding of how the brain processes information underlying the generation of behavior," he says.

**More information:** Oliver Skocek et al, High-speed volumetric imaging of neuronal activity in freely moving rodents, *Nature Methods* (2018). [DOI: 10.1038/s41592-018-0008-0](https://doi.org/10.1038/s41592-018-0008-0)

Provided by Rockefeller University

Citation: A new way to watch brain activity in action (2018, May 7) retrieved 6 May 2024 from <https://medicalxpress.com/news/2018-05-brain-action.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--