

Brain scientists identify 'cross talk' between neurons that control touch in mice

May 29 2018



Credit: CC0 Public Domain

Scientists report they have uncovered a previously overlooked connection between neurons in two distinct areas of the mammalian brain. The neurons, they say, control the sense of touch, and their

experiments in mice offer insights into mapping brain circuitry that is responsible for normal and abnormal perception and movements linked to touch.

Results of the experiments are summarized in the May 29 issue of *Cell Reports*.

"Neuroscientists have much to learn about the circuitry that organizes [touch](#). Because the sense of touch plays a significant role in how we move around and understand the world, and affects the perception of pain, it's important to know what [neurons](#) connect to other neurons and what happens to signals between these impulse-transmitting cells in the [brain](#)," says Daniel O'Connor, Ph.D., an associate professor of neuroscience at the Johns Hopkins University School of Medicine.

Often, he says, we experience touch passively, for example the feeling of clothes on our body or the feel of a chair we're sitting in. But when we actively touch, say by pressing down on an object to see if it's hard or moving our hand back and forth to see if it's smooth, we combine the sense of touch with [movement](#). And this processing—so vital to our lives and even survival—happens fast, sometimes in less than a thousandth of a second, says O'Connor.

In humans, O'Connor says, the reaction occurs in an area of the brain called the [somatosensory cortex](#), which runs along the area where a headband for earphones sits on the head. This strip of the brain is where all sensory input about touch is processed. It's divided into two main areas: S1, the primary somatosensory cortex, which runs along the headband region, and S2, the secondary somatosensory cortex, which is closer to the ears.

The S1 and S2 areas of the cortex are further divided into six layers of tissue. Working with [mice](#), O'Connor and his team undertook a close

look at the neurons in layer 4 of the cortex. It's long been thought that these layer 4 neurons in the cortex receive sensory input from the outside world and then broadcast it to other neurons in the same local region. Neurons send chemical and electric signals by extending arms, or "axons," to other neurons.

For the first part of the study, O'Connor and his team at the Johns Hopkins Solomon H. Snyder Department of Neuroscience, Brain Science Institute and Kavli Neuroscience Discovery Institute studied more than 47 mice by repeatedly touching their whiskers and shining a light through their skull. O'Connor says mouse and human brains have similar features, including cellular structures. Mice use their whiskers like humans use their fingers and hands, so many scientists focus their experiments on these hairlike structures in the animals as surrogates for human touch sensors.

To pinpoint the location of neurons responding to touch, the researchers shined light on the brain and used a quirk of brain physiology: the more brain activity, the more light is absorbed. Then, in the brain areas identified by the light experiment, the scientists injected chemical tracers that traveled along the neurons to find where they extended their axons and emitted chemical signals.

"To our surprise, we found in mice that layer 4 neurons in the S2 region were extending their axons beyond their local area, all the way to neurons in the S1 region," says O'Connor.

The S1 and S2 regions of the cortex in mice and humans sit alongside areas of the brain that control movement. "That's not a coincidence. We think it's by design, because there is a link between movement and sensation," says O'Connor.

O'Connor and his team expanded their research to explore how layer 4

neurons function. Among the mice, they monitored thousands of neurons—including layer 4 neurons that sent their axons to S1—with electrodes that record activity by the millisecond. The scientists wiggled whiskers on the mice with calibrated, computer-controlled movements.

The scientists were able to identify layer 4 neurons in both S1 and S2 cortical regions that were activated in the first 10 milliseconds of each whisker movement. "Because they react so fast, it implies that the feedback loop is happening as soon as the touch experience happens," says O'Connor.

Using sophisticated microscopes, the scientists then zoomed in on axons that extended from the S2 to S1 region. They moved the whiskers of the mice up and down and forward and backward, simulating how mice use their whiskers to perceive the direction of touch. About 10 percent of the layer 4 neurons were activated by specific directions of movement.

When O'Connor and his team blocked activity in layer 4 neurons in the S2 region using light-activated silencing molecules, neurons in the S1 region responded less specifically to up and down or back and forth movements of the whiskers—the neurons no longer "cared" much about the direction of movement.

"This tells us that the feedback loop between S2 layer 4 and S1 neurons improves encoding of information about the orientation or direction of touch," says O'Connor.

In addition to helping scientists understand the brain circuitry involved in touch, the work, says O'Connor, could one day inform the understanding and potentially treatment of some brain disorders involving movement and touch, such as strokes and chronic pain.

Provided by Johns Hopkins University School of Medicine

Citation: Brain scientists identify 'cross talk' between neurons that control touch in mice (2018, May 29) retrieved 26 April 2024 from <https://medicalxpress.com/news/2018-05-brain-scientists-neurons-mice.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.