

Chronic medical conditions are common in women with urinary incontinence

May 11 2018

New research published in *BJU International* indicates that women with urinary incontinence often have other chronic conditions. The findings have important implications for prevention and treatment.

In an analysis of data from the 2005-06 to 2011-12 US National Health and Nutrition Examination Surveys on 3,800 women with urinary incontinence, only 11% of women had no other chronic conditions.

Four patterns of chronic conditions emerged with differences by urinary incontinence type and severity. Within three of the four clusters, the most prevalent [chronic conditions](#) linked with increased cardiovascular risk—such as hypertension and high cholesterol—were associated with increased urinary incontinence risk. Also, pulmonary disease—specifically asthma—affected all of the women in a single cluster.

"Our data provide new evidence of the relationship between [chronic medical conditions](#) and [urinary incontinence](#) burden in women.

Specifically, identification of women with a low burden of incontinence and modifiable conditions—such as hypertension, high cholesterol, and asthma—may inform future prevention and treatment efforts," said lead author Dr. Alayne Markland, of the University of Alabama at Birmingham.

More information: Alayne D. Markland et al, Cluster analysis of multiple chronic conditions associated with urinary incontinence among

women in the USA, *BJU International* (2018). [DOI: 10.1111/bju.14246](https://doi.org/10.1111/bju.14246)

Provided by Wiley

Citation: Chronic medical conditions are common in women with urinary incontinence (2018, May 11) retrieved 6 May 2024 from <https://medicalxpress.com/news/2018-05-chronic-medical-conditions-common-women.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.