


Clinical trial to determine therapies for eradication of multiple myeloma

May 10 2018, by Savannah Koplon



Multiple myeloma is the second most common type of blood cancer in the U.S., impacting more than 25,000 people each year and disproportionately affecting African-American men. Credit: University of Alabama at Birmingham

Multiple myeloma is the second most common type of blood cancer in

the United States, impacting more than 25,000 people each year and disproportionately affecting African-American men. While patients have access to more quality treatment therapies than ever before, the disease is still regarded as an incurable condition that often leads to death for those affected.

However, researchers at the University of Alabama at Birmingham Division of Hematology and Oncology are leading and currently recruiting for a Phase II clinical trial that intends to provide newly diagnosed [multiple myeloma patients](#) an innovative [treatment](#) plan. The hope is that the therapies will eradicate the disease in a significant proportion of patients and measure disease response more accurately than ever before.

The clinical trial Monoclonal Antibody Sequential Therapy for Deep Remission in Multiple Myeloma, also known as MASTER, utilizes next generation sequencing technology to detect minimal residual disease down to a level of one cancer cell in 100,000—or 100- to 1,000-fold more sensitive than traditional methods to evaluate response. With a goal of enrolling 82 patients, those in the trial will be treated with a combination of anti-[myeloma](#) agents and immunotherapy that have a proven record of eliminating minimal residual disease, including the drug carfilzomib and the monoclonal antibody daratumumab, agents that are currently only approved to treat patients whose disease has returned.

"I believe for the first time that we have treatments that are effective enough to make it possible to eradicate multiple myeloma definitively in a substantial proportion of patients, along with having the technology to detect that the disease has been targeted and that treatment can be stopped," said Luciano Costa, M.D., Ph.D., scientist at the UAB Comprehensive Cancer Center, lead of the Hematologic-Malignancy Working Group, and principal investigator of the MASTER study. "That is what patients want, after all—a treatment that gives them the

possibility of eliminating any trace of the myeloma without having to be on therapy for the rest of their lives. It is a bold move, but bold moves are what our patients deserve."

This is one of the first [trials](#) in multiple myeloma to use minimal residual disease as primary endpoint and the very first one to modify therapy based on achievement of minimal residual disease eradication.

Up until now, existing therapies have been developed on trials that assigned treatment for a defined duration of time, irrespective of depth or speed of response, often followed by maintenance therapy for indefinite duration and with potentially adverse impact on risk of complications and cost. As it stands, patients with billions of cancer cells in their bodies and patients with true disease eradication will both appear to be in complete remission using current methods to measure disease response to therapy.

More than improving the results of the initial treatment, the MASTER study will treat patients for the necessary time to confirm elimination of [minimal residual disease](#) and then discontinue [therapy](#). Patients will be monitored for relapse at the molecular level, before [disease](#) becomes again symptomatic.

Provided by University of Alabama at Birmingham

Citation: Clinical trial to determine therapies for eradication of multiple myeloma (2018, May 10) retrieved 8 May 2024 from <https://medicalxpress.com/news/2018-05-clinical-trial-therapies-eradication-multiple.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--