


Providing clinicians feedback on their opioid prescribing data alters future prescribing

May 7 2018



Initial self-assessment, then information on prescribing patterns within peer distribution. Credit: Kirsty Challen, B.Sc., MBCHB, MRES, PH.D., Lancashire Teaching Hospitals, United Kingdom.

Asking emergency department (ED) providers to self-identify their opioid prescribing practices and then providing them with timely, clinically relevant, individualized, and actionable feedback on their actual opioid prescribing data, significantly decreases future opioid prescribing among providers who underestimate their baseline prescribing. That is the finding of a study to be published in the May 2018 issue of *Academic Emergency Medicine* (AEM), a journal of the Society for Academic Emergency Medicine (SAEM).

The lead author of the study is Sean S. Michael, MD, assistant professor of [emergency medicine](#) at the University of Colorado Denver School of Medicine.

The study, by Michael, et al, found that providers who initially underestimated opiate prescription rates had statistically significant decreases in opiate prescribing when compared to providers who were more accurate in initial self-assessment of prescribing rates. The randomized trial exposes important gaps in providers' self-perceptions of opioid prescribing and demonstrates that a simple, data-driven intervention using query-reveal methodology may decrease future prescribing, particularly among providers who underestimate their own prescribing practices.

The findings suggest that current and future interventions in which provider adherence to guidelines is expected, including opioid interventions outside the emergency department and other quality and safety initiatives, should directly address the potential barrier of inaccurate provider self-awareness.

Gail D'Onofrio, MD, MS, professor and chair, Yale School of Medicine Department of Emergency Medicine, who is internationally known for her work in developing and testing interventions for department patients with unhealthy alcohol and other drug use, commented:

"The results have real world applicability. A simple intervention enhancing safe prescribing is one of several opportunities the [emergency](#) physician has to combat the current opioid epidemic."

More information: Sean S. Michael et al, Effect of a Data-driven Intervention on Opioid Prescribing Intensity Among Emergency Department Providers: A Randomized Controlled Trial, *Academic Emergency Medicine* (2018). [DOI: 10.1111/acem.13400](https://doi.org/10.1111/acem.13400)

Provided by Society for Academic Emergency Medicine

Citation: Providing clinicians feedback on their opioid prescribing data alters future prescribing (2018, May 7) retrieved 5 May 2024 from <https://medicalxpress.com/news/2018-05-clinicians-feedback-opioid-future.html>

| |
|--|
| <p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p> |
|--|