

# Records point to drug-drug interaction

May 8 2018

---

Daptomycin is an antibiotic used to treat serious bacterial infections. Statins are drugs prescribed on a long-term basis to prevent or reduce cardiovascular disease by lowering LDL-cholesterol.

In a retrospective study, researchers at Vanderbilt University Medical Center found that VUMC patients who continued taking a [statin](#) while taking daptomycin had 2.6 times greater risk of developing myopathy ([muscle weakness](#)) and 4.67 times greater risk of a more severe form of [muscle damage](#) called rhabdomyolysis.

"This is the first study to provide strong evidence supporting this association," reported Ryan Dare, MD, MS, and co-authors in the journal *Clinical Infectious Diseases*. Echoing the maker of daptomycin, the study authors recommend that doctors consider withholding statins during daptomycin therapy.

Among patients on statins seen at VUMC between 2004-2015, the researchers found 128 cases of daptomycin-associated myopathy, including 25 cases of rhabdomyolysis; they matched these patients to myopathy-free controls, adjusting their analyses to account for a range of risk factors.

**More information:** Ryan K Dare et al. Statin Coadministration Increases the Risk of Daptomycin-Associated Myopathy, *Clinical Infectious Diseases* (2018). [DOI: 10.1093/cid/ciy287](https://doi.org/10.1093/cid/ciy287)

Provided by Vanderbilt University

Citation: Records point to drug-drug interaction (2018, May 8) retrieved 20 April 2024 from <https://medicalxpress.com/news/2018-05-drug-drug-interaction.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.