

Eggs not linked to cardiovascular risk, despite conflicting advice

May 7 2018

University of Sydney researchers aim to help clear up conflicting dietary advice around egg consumption, as a new study finds eating up to 12 eggs per week for a year did not increase cardiovascular risk factors in people with pre-diabetes and type 2 diabetes.

Published in the *American Journal of Clinical Nutrition* today, the research extends on a previous study that found similar results over a period of three months.

Led by Dr Nick Fuller from the University's Boden Institute of Obesity, Nutrition, Exercise and Eating Disorders at the Charles Perkins Centre, the research was conducted with the University of Sydney's Sydney Medical School and the Royal Prince Alfred Hospital.

In the initial trial, participants aimed to maintain their [weight](#) while embarking on a high-egg (12 eggs per week) or low-egg (less than two eggs per week) [diet](#), with no difference in [cardiovascular risk](#) markers identified at the end of three months.

The same participants then embarked on a [weight loss](#) diet for an additional three months, while continuing their high or low egg consumption. For a further six months - up to 12 [months](#) in total - participants were followed up by researchers and continued their high or low egg intake.

At all stages, both groups showed no adverse changes in cardiovascular

risk markers and achieved equivalent weight loss - regardless of their level of egg consumption, Dr Fuller explained.

"Despite differing advice around safe levels of egg consumption for people with pre-diabetes and type 2 diabetes, our research indicates people do not need to hold back from eating eggs if this is part of a healthy diet," Dr Fuller said.

"A healthy diet as prescribed in this study emphasised replacing saturated fats (such as butter) with monounsaturated and polyunsaturated fats (such as avocado and olive oil)," he added.

The extended study tracked a broad range of [cardiovascular risk factors](#) including cholesterol, blood sugar and blood pressure, with no significant difference in results between the high egg and low egg groups.

"While eggs themselves are high in dietary cholesterol - and people with type 2 diabetes tend to have higher levels of the 'bad' low density lipoprotein (LDL) cholesterol - this study supports existing research that shows consumption of eggs has little effect on the levels of cholesterol in the blood of the people eating them," Dr Fuller explained.

Dr Fuller said the findings of the study were important due to the potential health benefits of [eggs](#) for people with pre-diabetes and type 2 diabetes, as well as the general population.

"Eggs are a source of protein and micronutrients that could support a range of health and dietary factors including helping to regulate the intake of fat and carbohydrate, eye and heart health, healthy blood vessels and healthy pregnancies."

The different egg diets also appeared to have no impact on weight, Dr Fuller said.

"Interestingly, people on both the high egg and low egg diets lost an equivalent amount of weight - and continued to lose weight after the three month intended weight loss phase had ended," he said.

More information: Nicholas R Fuller et al, Effect of a high-egg diet on cardiometabolic risk factors in people with type 2 diabetes: the Diabetes and Egg (DIABEGG) Study—randomized weight-loss and follow-up phase, *The American Journal of Clinical Nutrition* (2018). [DOI: 10.1093/ajcn/nqy048](https://doi.org/10.1093/ajcn/nqy048)

Provided by University of Sydney

Citation: Eggs not linked to cardiovascular risk, despite conflicting advice (2018, May 7)
retrieved 27 April 2024 from
<https://medicalxpress.com/news/2018-05-eggs-linked-cardiovascular-conflicting-advice.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.