

Food safety: Are the sniff test, the fivesecond rule and rare burgers safe?

May 10 2018, by Cathy Moir

If you drop your food, it's less about time spent on the floor and more about the surfaces. Credit: Mai Lam/The Conversation NY-BD-CC, CC BY-SA

There are many rules in food safety lore, some that have a basis in fact, and some that are purely grounded in convenience. But it's important to look at the evidence to see which category common rules fall under.

1. The 'sniff' test



Often when a <u>food</u> has spoiled, it will smell bad. This leads many to believe "no stench = OK to eat". But this isn't always the case. The <u>microorganisms</u> (bacteria, yeasts and moulds) that spoil food by making it smelly, slimy or mouldy might not give you food poisoning.

But pathogenic (disease-causing) bacteria, such as salmonella, campylobacter, E.coli and listeria, which do make people sick, don't always cause obvious changes in food when they grow. Sometimes simply being present at low numbers and then consumed is enough to result in illness.

Having said that, this isn't an invitation to consume obviously spoiled food. Spoilage is a good indicator food has been left too long and "bad" microorganisms, including pathogens, may also have grown.

In order to steer clear of nasty bugs in food, observe "use by" dates, refrigerate foods that need to be kept cold (this slows down the microbes), cook foods properly (this kills the microbes) and prevent contact and cross contamination between ready-to-eat foods such as salads, with raw food such as <u>meat</u> that still needs to be cooked.

2. The 'five second' rule

Whether it's one, three, five seconds or some other number, we've all heard some version of this call when someone has dropped food on the floor. But is it true harmful bacteria need a few seconds to hitch a ride on your dropped slice of pizza?

In one <u>peer-reviewed study</u>, four food types were tested (watermelon, bread, bread and butter, and gummy sweets) with four different surfaces (<u>stainless steel</u>, ceramic tile, wood and carpet) that were contaminated with bacteria. Contact time, food type and surface all significantly affected the amount of contamination that occurred. The study found:





Biscuits are dry and thus not super attractive to bacteria. Credit: The Creative Exchange/Unsplash

- time is not necessarily of the essence as microorganisms from one surface can instantaneously contaminate another. But it's true the longer contact time the more contamination can occur
- higher moisture foods (such as watermelon) allowed transfer of more microorganisms compared to the other foods. The gummy sweets, which are likely to have the driest surface, showed the weakest transfer of bacteria from the contact surface



• the weakest microbial transfer occurred when food was dropped on to carpet compared to stainless steel, and tile in particular. The authors hypothesised bacteria attaches better to carpet as it's more absorbent, meaning it's less likely to transfer to the food.

While it's true dropped food can become contaminated with microorganisms from the floor or environment, the majority of those microorganisms in a normal home are likely to be harmless to human health.

3. Rare meat

When cooking and reheating meat, there are some simple rules to follow. Whole pieces of meat muscle such as steak, pork and lamb can be cooked on the outside, say barbequed or pan fried, so they're still rare on the inside.

Historically, under-cooked pork has been feared due to a parasitic worm, but this has never been seen in Australian pigs.

Poultry and all minced, rolled, stuffed, tenderised and similar types of meat (including burgers) need to be cooked right through. This difference relates to where microorganisms are found on the meat.

We know microorganisms live on the surface of raw meat because animals naturally harbour microorganisms. That's why just cooking the surface of a whole piece of muscle meat is sufficient (excluding poultry), because that will kill any potentially <u>harmful bacteria</u>.

When that meat is minced, rolled, stuffed, mechanically tenderised or turned into patties or sausages, the surface of the meat and what it's carrying is then mixed through the whole product. It's also possible for chicken tissue to be colonised by <u>bacteria</u> (which just doesn't happen



with other animal meat types). That's why these types of meat products need to be cooked through to the centre.

The best way to tell if meat is cooked is to use a meat thermometer. These can be purchased from homeware and hardware stores. Poultry and minced, rolled, stuffed, tenderised meats need to be cooked right through and to a temperature of 75°C. Insert the thermometer into the thickest part of the meat. If you don't have a thermometer, check the juices run clear and not pink.

This article was originally published on <u>The Conversation</u>. Read the <u>original article</u>.

Provided by The Conversation

Citation: Food safety: Are the sniff test, the five-second rule and rare burgers safe? (2018, May 10) retrieved 15 May 2024 from https://medicalxpress.com/news/2018-05-food-safety-five-second-rare-burgers.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.