

# New guidelines to help clinicians manage GSM in women with breast cancer

May 17 2018

---

Although the genitourinary syndrome of menopause (GSM) is more prevalent in survivors of breast cancer than in other menopausal women, it is commonly undiagnosed and untreated. This led The North American Menopause Society (NAMS) and The International Society for the Study of Women's Sexual Health (ISSWSH) to form a multidisciplinary Consensus Panel to develop recommendations for clinicians to manage GSM in women with or at high risk for breast cancer. The recommendations are published online in NAMS' official journal, *Menopause*.

For survivors of breast cancer, not only is GSM more prevalent, but these women may experience its symptoms earlier because of cancer treatments. With the lack of data regarding the safety of vaginal hormone treatments and other alternatives for women with or at [high risk](#) for breast cancer, many have avoided using hormone treatments, and [clinicians](#) have often been reluctant to prescribe them. Left untreated, GSM can potentially adversely affect quality of life and partner relationships because of ongoing symptoms, including vulvovaginal dryness, burning, or irritation; painful intercourse; and recurrent urinary tract infections. It is estimated that more than 2 million survivors of [breast cancer](#) in the United States are affected by GSM, with most receiving no treatment.

The Consensus Panel of 16 experts created the consensus recommendations after a comprehensive, evidence-based review of existing literature. In addition, the Consensus Panel agreed that decision

making regarding the type of treatment for GSM should be individualized and include balancing the risk of [cancer](#) recurrence with the severity of symptoms and effect on quality of life. The recommendations offer clinicians caring for this population of women some guidance regarding therapeutic decision making in the absence of [clinical trial data](#).

"We want clinicians caring for this at-risk population to be armed with as much information as possible so that they can provide the best course of treatment for their patients," says Dr. JoAnn Pinkerton, executive director of NAMS. "Until additional studies are undertaken in this area, we are hopeful these consensus recommendations will provide added confidence for clinicians to move forward with treatment options that will provide relief to [women](#) from GSM symptoms."

Provided by The North American Menopause Society

Citation: New guidelines to help clinicians manage GSM in women with breast cancer (2018, May 17) retrieved 20 April 2024 from <https://medicalxpress.com/news/2018-05-guidelines-clinicians-gsm-women-breast.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--