

# Hospitalized patients with acute kidney injury have increased risk of heart failure

May 18 2018

---

Hospitalized patients who experience acute kidney injury face a 44 percent greater risk of heart failure during their first year after leaving the hospital, according to new Kaiser Permanente research published today in the *Clinical Journal of the American Society of Nephrology*.

Acute [kidney](#) injury is a sudden decline in the ability of the kidneys to perform their normal function of filtering waste products from blood. People who are hospitalized for a wide variety of conditions, from major surgery to severe infection, are at risk for acute kidney injury, which can increase a patient's long-term risk of death even after they are successfully treated and sent home.

"While Kaiser Permanente has had success in lowering rates of acute kidney injury in hospitalized [patients](#), rates are rising nationwide, and even minor injury is linked to a higher risk of death," said Alan S. Go, MD, director of the Comprehensive Clinical Research Unit within the Kaiser Permanente Northern California Division of Research and lead author of the new study.

Earlier research by Dr. Go and colleagues linked hospitalized acute kidney injury to elevated risk of [high blood pressure](#). In this new study, which adds to that body of work, the research team examined [electronic health records](#) of patients who were admitted to one of Kaiser Permanente's 21 hospitals in Northern California at some point between 2006 and 2013. They investigated whether the risk of various [cardiovascular events](#) within one year of leaving the hospital differed

between patients who experienced acute kidney injury while hospitalized and those who did not.

Out of 146,931 hospitalized patients included in the analysis, 31,245 experienced acute kidney injury. To help ensure an accurate comparison, patients with acute kidney injury were statistically matched to patients without acute kidney injury, according to similar demographics, length of hospital stay, medications they took, how acutely ill they were and other characteristics.

Roughly 1 in 25 people who are hospitalized for any reason will experience heart [failure](#) within a year. The analysis found that even after accounting for remaining differences, patients who have an acute kidney injury during hospitalization experienced a 44 percent higher relative risk of heart failure within one year of discharge, compared to those without acute kidney injury. Rates of other cardiovascular events linked to atherosclerosis (narrowing of the arteries) did not differ significantly between the two groups.

"The number of atherosclerotic events in patients with acute kidney injury was lower than we expected," Dr. Go said. "Overall, our results highlight heart failure as a key risk for patients who experienced acute kidney injury in the hospital."

According to senior author Kathleen D. Liu, MD, Ph.D., of the University of California, San Francisco, the new findings suggest the need for doctors to be more vigilant in looking for signs of heart failure in patients who experienced acute kidney [injury](#) during hospitalization. "Earlier detection of heart failure symptoms in these patients could potentially save lives."

Meanwhile, Dr. Go and others are examining the mechanisms underlying the increased risk. "Kidney damage affects a number of biological

pathways, including inflammation and mineral metabolism," Dr. Go said. "If we could understand how these changes increase risk of [heart](#) failure, we may be able to develop new strategies for prevention."

Provided by Kaiser Permanente

Citation: Hospitalized patients with acute kidney injury have increased risk of heart failure (2018, May 18) retrieved 26 April 2024 from <https://medicalxpress.com/news/2018-05-hospitalized-patients-acute-kidney-injury.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--