


Label-free method for rapid cancer diagnosis

May 21 2018



Claus Kuepper, Frederik Großerueschkamp, Angela Kallenbach-Thieltges and Klaus Gerwert (left to right) developed novel methods of cancer diagnostics.
Credit: RUB, Marquard

Researchers at Ruhr-Universität Bochum have deployed a novel infrared (IR) microscope with quantum cascade lasers in order to analyse tissue samples taken during routine clinical procedures for colorectal cancer

diagnosis. The Fourier transform infrared (FTIR) microscope used today had not yet been established as a diagnostic tool in hospitals, as the analyses previously took too long. By utilising the new laser technology, the researchers reduced the time required for analysis from one day to a few minutes. Coupled with bioinformatic image analysis, the IR microscope performs label-free classification of cancer tissue and can be fully automated.

Bochum-based researchers led by Prof Dr. Klaus Gerwert, Dr. Angela Kallenbach-Thieltges, Dr. Frederik Großerüschkamp and Claus Küpper from the Department of Biophysics published a report on the project in *Scientific Reports*.

In previous studies, the biophysicists had already demonstrated the potential OF THE the FTIR microscope in combination with FTIR imaging as a diagnostic tool for the classification of tissue. Unlike traditional clinical rapid diagnostic tests, which take approximately 20 minutes, FTIR imaging used to take a whole day. Now, the researchers have considerably simplified the measurement set-up and replaced FT technology with quantum cascade laser technology, akin to replacing a weak light bulb that emits diffuse light with precisely bundled, intense laser light.

In collaboration with the Institute of Pathology at Ruhr-Universität, headed by Prof Dr. Andrea Tannapfel, they used IR imaging to analyse [120 tissue samples](#) taken from patients suffering from [colorectal cancer](#). The analysis was based on algorithms developed by the biophysicists in-house that were used for colouring the IR images of tissue samples at the computer. The results corresponded to 97 percent with traditional histopathological analyses. "We have thus reduced the measurement period by a factor of 160," describes Frederik Großerüschkamp.

As control, the measurements were carried out using two pieces of

equipment, and the analyses were performed by several users; this did not affect the results. "The method is now very fast, reliable and does not depend on a specific device or a specific user," says Angela Kallenbach-Thieltges. "This opens up new avenues for automated classification of [tissue](#) samples taken directly from the patient."

In future, the team intends to incorporate the method into clinical workflow. "Automated image analysis might be deployed as a time-saving [diagnostic tool](#), which might possibly even be used in situ," anticipates pathologist Andrea Tannapfel.

Colorectal cancer is one of the most common tumour diseases, and it is very treatable following early diagnosis. "The results of the study give rise to hope that highly precise therapy is within reach, which can be personalised for each individual patient and, consequently, will ultimately prove more successful than traditional approaches," says Klaus Gerwert.

More information: Claus Kuepper et al, Quantum Cascade Laser-Based Infrared Microscopy for Label-Free and Automated Cancer Classification in Tissue Sections, *Scientific Reports* (2018). [DOI: 10.1038/s41598-018-26098-w](#)

Provided by Ruhr-Universitaet-Bochum

Citation: Label-free method for rapid cancer diagnosis (2018, May 21) retrieved 19 April 2024 from <https://medicalxpress.com/news/2018-05-label-free-method-rapid-cancer-diagnosis.html>

| |
|--|
| <p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p> |
|--|