

Many patients with rectal cancer may not need chemoradiotherapy

May 11 2018

It's unclear whether all patients with advanced rectal cancer need chemoradiotherapy, or whether some can forego the treatment and therefore be spared its side effects. A new *BJS* (*British Journal of Surgery*) study found that patients who are not at high risk of local recurrence can safely undergo surgery without receiving chemoradiotherapy. Investigators estimate that this strategy could reduce the number of patients who receive chemoradiotherapy by roughly half, with only a minimal risk of undertreatment.

In the observational trial, 545 patients with cancer of the rectum were stratified into low-and high-risk groups. Chemoradiotherapy was administered to patients at <u>high risk</u> and was avoided in 42% of patients. The 3- and 5-year local recurrence rates of cancer were similarly low in patients who had surgery alone (a procedure called total mesorectal excision) versus <u>patients</u> who received chemoradiotherapy followed by surgery.

"The results are an important step to an individualized therapy of rectal <u>cancer</u>," said co-author Prof. Theodor Junginger, of the University Medical Centre Mainz, in Germany. "The concept avoids adverse effects of chemoradiotherapy, shortens duration of treatment, and saves costs."

More information: R. Ruppert et al, Oncological outcome after MRIbased selection for neoadjuvant chemoradiotherapy in the OCUM Rectal Cancer Trial, *British Journal of Surgery* (2018). DOI: 10.1002/bjs.10879



Provided by Wiley

Citation: Many patients with rectal cancer may not need chemoradiotherapy (2018, May 11) retrieved 5 May 2024 from https://medicalxpress.com/news/2018-05-patients-rectal-cancer-chemoradiotherapy.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.