

# Should patients in remission stop taking expensive rheumatoid arthritis drugs?

May 11 2018

---

Stopping expensive biological drugs used to treat rheumatoid arthritis in patients who are in remission or who have low disease activity can save considerable costs, but it results in a small loss of quality-adjusted life years, according to a recent *Arthritis & Rheumatology* study.

In the study, stopping [tumor necrosis factor](#) inhibitors in patients with stable low disease activity, on average, was associated with a cost saving of €7,133, a loss of 0.022 quality-adjusted life years, and an increase of 0.41 arthritis flares per patient per year.

"The subpopulation of patients receiving biological disease modifying anti-rheumatic drugs including tumor necrosis factor inhibitors... has increased over time and accounted for up to 20% of the population of [rheumatoid arthritis](#) patients in various Western healthcare systems," the authors wrote.

**More information:** An Tran-Duy et al, An Economic Evaluation of Stopping versus Continuing TNF-Inhibitor Treatment in Rheumatoid Arthritis Patients in Remission or Low Disease Activity: results from the POET randomized trial, *Arthritis & Rheumatology* (2018). [DOI: 10.1002/art.40546](#)

Provided by Wiley

Citation: Should patients in remission stop taking expensive rheumatoid arthritis drugs? (2018, May 11) retrieved 18 April 2024 from <https://medicalxpress.com/news/2018-05-patients-remission-expensive-rheumatoid-arthritis.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.