

# It's OK to skip your period while on the pill

May 17 2018, by Mia Schaumberg

---



Women commonly suppress their period around special events and holidays.  
Credit: Haley Phelps

The pill is the most common form of contraception in Australia, with [up to 60%](#) of young women choosing this option to prevent pregnancy.

Women on the [pill](#) are able to manipulate or suppress their [menstrual](#)

[cycles](#) to have fewer "periods," or to avoid bleeding at important or inconvenient times. But it's best to consult your GP to work out a plan that suits you and be on the lookout for any side effects.

## How does the pill work?

The combined [oral contraceptive pill](#) (known as "the pill") provides a daily dose of synthetic ovarian hormones oestradiol and progestogen, which [suppress the natural production](#) of ovarian hormones (oestrogen and progesterone) to prevent ovulation.

The idea of an oral contraceptive using ovarian hormones was [first proposed](#) in the 1950s by American researcher Dr. Gregory Pincus. His idea was to mimic the natural cycle, with 21 days of hormone, followed by seven days of no [hormone](#) to allow for a "withdrawal bleed."

While the withdrawal bleed may seem like a period, it has little physiological relation to menstruation and may be lighter, shorter and less painful.

Nonetheless, [early research described](#) the 21/7 regimen as a "morally permissible variant of the rhythm method," hopefully leading to acceptance by women, their doctors, and the Catholic Church.

## How did menstrual suppression come about?

In 1977, the [first study](#) was published on an extended oral contraceptive regimen of 84/7 (84 days of active pills and seven days of no hormones). This showed reducing the frequency of menstruation to every three months was safe.

Since then, researchers have studied the various ways the pill can be used

to suppress menstruation.

Most studies show the practice is well-accepted by patients. An [84-day cycle is usually recommended](#), allowing four withdrawal bleeds per year. This schedule minimises the risk of [endometrial hyperplasia](#), an abnormal thickening of the lining of the uterus, which is also a risk factor for endometrial cancers.

While the most common way to manipulate the menstrual cycle is through extending the days of oral contraceptive use, there are several other [long-acting reversible hormone contraceptive methods](#) that can be used to suppression menstruation for months or years. These include Implanon (a hormonal implant in the arm), Mirena (a device that sits in the uterus), or Depo-provera (an injection).



The sugar pills don't have any active ingredients. Credit: [Thought Catalog](#)

## How common is menstrual suppression?

In my [recent survey of young Australian women](#), 74% reported manipulating their menstrual cycle at least once in the previous year. Almost one-third of respondents had manipulated their cycle two to three times. But less than 1% had used the pill to stop menstruation altogether.

The most commonly cited reason was to be able to suppress a period around special events or holidays, with 75% of the women rating this as important or very important.

Suppressing menstruation can be more practical than just ensuring you don't have your period for a wedding or important exam. In a [survey of deployed US female aviation personnel](#), 60% chose to suppress [menstruation](#) to ensure they could meet the demands of their job.

[More than 50% of competitive athletes](#) use the pill to ensure they don't have their period on competition day.

What about women who suffer severe or painful menstrual symptoms, like heavy bleeding or endometriosis?

Menstrual manipulation or suppression can also help women with endometriosis, heavy bleeding or painful periods to minimise menstrual symptoms. Currently, [about 30% of women](#) take four or more sick days per year due to premenstrual symptoms alone. Those who suffer severe

symptoms may be absent from work or school for one day every month. This is a large potential economic burden.

Menstrual manipulation is a strategy to reduce absenteeism from work and school and may be of benefit at both an individual and societal level.

## **This all sounds so good – so what are the risks?**

Some arguments suggest that by skipping the "sugar pills," [women](#) are effectively increasing the overall exposure to synthetic hormones by up to 25%. This may lead to a greater risk of depression, blood clots, or oestrogenic cancers.

Unfortunately, there is no research yet to support or refute these claims.

However there is evidence that menstrual manipulation can lead to greater incidence of "spot" or "breakthrough" bleeding. Women can also find it more difficult to tell if they become pregnant.

While there is no evidence to suggest menstrual manipulation or suppression is harmful, it's a good idea to discuss it with your GP or gynaecologist, who can assess your individual circumstances and potential risk factors.

Under medical guidance, you can decide on a system to suppress your periods, and have a plan to contact your doctor if you experience any side effects such as abdominal pain, chest pain, bleeding, vision changes or severe leg pain.

This article was originally published on [The Conversation](#). Read the [original article](#).

Provided by The Conversation

Citation: It's OK to skip your period while on the pill (2018, May 17) retrieved 6 May 2024 from <https://medicalxpress.com/news/2018-05-period-pill.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.