



Post-mortem CT angiography illuminates causes of death

May 1 2018



a.



b.

Graph of additional postmortem findings. (a) Additional findings obtained by using imaging with autopsy. Autopsy findings as a percentage of all findings are in red. Additional findings not observed at autopsy but identified at postmortem CT are in blue. Green indicates findings undiscovered at both autopsy and postmortem CT but detected at postmortem CT angiography. Credit: Radiological Society of North America



CT angiography is a useful adjunct to autopsy that is likely to increase the quality of post-mortem diagnosis, according to a study appearing online in the journal *Radiology*. Researchers said the findings could have important implications for criminal investigations and hospital quality control efforts.

High-quality post-mortem investigations are vital for a number of reasons; most notably, as evidence in criminal investigations for which the body represents the main proof of a crime, and as a quality control measure for surgical interventions and medical treatment at hospitals.

Modern imaging techniques like CT and MRI are often used in forensic pathology. However, post-mortem CT is limited by relatively low soft tissue contrast and poor visualization of the vascular system. A technique known as post-mortem CT angiography, or PMCTA, was developed to address those limitations by introducing contrast agent into the body.




"The idea of the PMCTA is to simulate a post-mortem circulation by establishing a post-mortem perfusion of the vascular system," said study lead author Silke Grabherr, M.D., Ph.D., from the University Center of Legal Medicine Lausanne-Geneva in Lausanne, Switzerland. "Therefore, the heart is 'replaced' by a perfusion device, similar to the heart-lung

machine that replaces the heart during heart surgery."



a.

b.



c.

d.

e.


(a) Axial cervical postmortem CT scan and (b) zoomed section of the cricoid cartilage (box in a) of a 27-year-old woman who died of strangulation. Three-dimensional volumerendered reconstructions from cranial (c), left lateral oblique (d), and right lateral oblique (e) views. Credit: Radiological Society of North America

To assess the performance of PMCTA, Dr. Grabherr and colleagues at nine centers in Europe performed conventional autopsies on 500 human bodies that had undergone PMCTA. All CT images were read by an experienced team including one forensic pathologist and one radiologist who were blinded to the autopsy findings.

Autopsy was performed on the day of the PMCTA, or the following day, in all cases. After examination of 500 bodies, a total of 18,654 findings were recorded.

PMCTA performed surprisingly well compared to conventional autopsy. Out of 18,654 total findings, PMCTA found 90 percent, compared with 61 percent for autopsies. PMCTA missed only 10 percent of forensically essential findings, while autopsy missed 23 percent.

"This method could—in many cases—be an alternative to invasive autopsy if such an opening of the body is not possible," Dr. Grabherr said. "This offers new investigation possibilities, for example, in countries where a conventional autopsy is not accepted or in cases where family members can refuse it."



a.



b.



c.



d.

(a) Postmortem CT and (b-d) arterial phase postmortem CT angiography images in a 59-year-old woman who died of internal exsanguination shortly after Whipple surgery. Credit: Radiological Society of North America

PMCTA was significantly superior to autopsy at identifying skeletal and vascular lesions, or areas of damage in the bones and blood vessels. PMCTA identified 96 percent of skeletal lesions and 94 percent of vascular lesions, compared with 65 percent for autopsy. These lesions can provide important information in post-mortem examinations, Dr. Grabherr said.

"The combination of bone lesions and vascular [lesions](#) is especially seen in cases of traumatic death, such as falls from height, traffic accidents, ballistic trauma and sharp trauma in homicides and suicides," she said. "This means that PMCTA is an excellent choice to investigate such cases and can be used combined with or even independently of conventional [autopsy](#) for investigating in these cases."

In the future, the research group plans to study the combination of post-mortem angiography with MRI in order to increase the sensitivity of findings related to organs like the brain and liver.

More information: "Postmortem CT Angiography Compared with Autopsy: A Forensic Multicenter Study" *Radiology*, 2018.

Provided by Radiological Society of North America

Citation: Post-mortem CT angiography illuminates causes of death (2018, May 1) retrieved 24 April 2024 from <https://medicalxpress.com/news/2018-05-post-mortem-ct-angiography-illuminates-death.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--