


Restricting fluids post abdominal surgery doesn't up survival

May 10 2018



(HealthDay)—A restrictive fluid regimen is not associated with

increased disability-free survival for patients at increased risk of complications during major abdominal surgery, but is associated with increased acute kidney injury, according to a study published online May 10 in the *New England Journal of Medicine*. The research was published to coincide with the annual meeting of the Australian and New Zealand College of Anaesthetists, held from May 7 to 11 in Sydney.

Paul S. Myles, M.P.H., D.Sc., from Monash University in Melbourne, Australia, and colleagues randomized 3,000 patients who had an increased risk of complications while undergoing major abdominal surgery to receive a restrictive or liberal intravenous-[fluid](#) regimen during and up to 24 hours after surgery (1,490 and 1,493 patients, respectively).

The researchers found that at one-year, the rate of disability-free survival was 81.9 and 82.3 percent in the restrictive and liberal fluid groups, respectively (hazard ratio for death or disability, 1.05; 95 percent confidence interval, 0.88 to 1.24; P = 0.61). The rate of [acute kidney injury](#) was 8.6 and 5.0 percent in the restrictive and liberal fluid groups, respectively (P

"Among [patients](#) at increased risk for complications during major abdominal [surgery](#), a restrictive fluid regimen was not associated with a higher rate of disability-free survival than a liberal fluid regimen and was associated with a higher rate of acute kidney injury," the authors write.

More information: [Abstract/Full Text](#)
[Editorial](#)
[More Information](#)

Copyright © 2018 [HealthDay](#). All rights reserved.

Citation: Restricting fluids post abdominal surgery doesn't up survival (2018, May 10) retrieved 25 April 2024 from

<https://medicalxpress.com/news/2018-05-restricting-fluids-abdominal-surgery-doesnt.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.