

Screening for colorectal cancer in under-55-year-olds with family history—benefit unclear

May 30 2018

Between the age of 50 and 54 years, all statutory health insurance fund members in Germany are entitled to immunological faecal occult blood testing and, in the case of a suspicious result, to a colonoscopy. From the age of 55 years onwards, members of the health insurance funds can choose between faecal occult blood testing every two years or two colonoscopies at least ten years apart.

The Federal Joint Committee (G-BA) is considering to abolish these fixed age limits for this [screening](#). As a result, testing could be offered earlier or more often in groups of [people](#) at risk (risk-adapted screening).

Focus on people under 55 years of age with a family history of colorectal cancer

The G-BA therefore commissioned the German Institute for Quality and Efficiency in Health Care (IQWiG) to update a report from the year 2013 in an accelerated process known as "rapid report" on the following question: Can people under 55 years of age with a family [history](#) of [colorectal cancer](#) benefit from a screening test?

The IQWiG researchers identified two further studies that are principally relevant for the research question. These studies provided no answers to the research question of the rapid report, however: Due to a lack of new findings, benefit or harm of the screening for people under

the age of 55 years with a family history remain unclear.

Family history of colorectal cancer increases risk

Based on the available studies, only the question regarding family risk could be sufficiently answered in IQWiG's final report from 2013: Under 55-year-olds with at least one first-degree relative with colorectal [cancer](#) have a 1.7 to 4.1 times higher risk of also developing colorectal cancer than people of the same age without a family history of colorectal cancer.

Two new studies on the comparison of screening procedures

The current search found two studies investigating whether colorectal cancer and advanced adenoma are less common in people with a [family](#) history of colorectal cancer if certain screening procedures are used: The FACTS study investigated whether it makes a difference if colonoscopy is repeated after three years or only after six years (surveillance interval). The COLONFAM study compared colonoscopy with immunological faecal occult blood testing. Due to a lack of conclusive results, these studies could not answer the study questions.

Benefit of risk-adapted screening remains unclear

Like the 2013 final report, the current rapid [report](#) therefore concluded that the benefit of screening for under 55-year-olds with a [family history](#) of colorectal cancer is unclear. A potential harm from screening tests also remains unclear: For instance, harm can be caused by people being wrongly allocated to the risk group, causing unnecessary psychological stress.

In view of this data situation, it must be carefully considered whether and how a risk-adapted screening strategy should be introduced.

More information: [www.iqwig.de/en/projects-resul... ion-s11-01.8014.html](http://www.iqwig.de/en/projects-results-ion-s11-01.8014.html)

Provided by Institute for Quality and Efficiency in Health Care

Citation: Screening for colorectal cancer in under-55-year-olds with family history—benefit unclear (2018, May 30) retrieved 25 April 2024 from <https://medicalxpress.com/news/2018-05-screening-colorectal-cancer-under-year-olds-family.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.