

Simple walking test helps predict risk for cognitive issues after heart surgery

May 10 2018

The distance a patient can walk in 6-minutes before a heart operation may be a clue to whether that patient will develop problems with memory, concentration, and attention after the procedure, according to a study published online today in *The Annals of Thoracic Surgery*.

Broadly speaking, a decline in cognitive performance after [surgery](#) is known as postoperative cognitive dysfunction (POCD). With POCD, a patient's mental aptitude is weaker after surgery, resulting not only in a greater risk of complications, but also a lesser quality of life. Cognitive deterioration is increasingly recognized as a common occurrence after major surgery, especially among older adult [patients](#).

"This study indicates that the easy and inexpensive 6-minute walk distance (6MWD) is a valuable assessment for identifying patients at a high risk for POCD," said Kazuhiro Hayashi, PT, MSc, of Nagoya University Hospital in Japan. "If we are able to identify patients who are at risk for POCD, we can provide early treatment and encourage them to better understand the dysfunction."

For this study, Hayashi and colleagues identified 181 patients who were undergoing non-emergency heart surgery between March 2014 and August 2015 at Nagoya University Hospital. The mean age of the patients was 71.4 years.

Patients performed the 6MWD test upon admission for their operations. Functional exercise capacity was measured by having patients walk the

length of a predetermined course at their own pace while attempting to cover as much ground as possible in 6 minutes. The distance covered in that duration was measured to the nearest meter. According to the results of this study, a low 6MWD was an associated risk factor for POCD after cardiac surgery. In fact, the lower the 6MWD was, the more significant the reduction in cognitive function postoperatively was. Of the study participants, 51 (28 percent) developed POCD.

"It is increasingly recognized that a patient's fitness level has an impact on how well he/she does after a surgical procedure," said Rakesh C. Arora, MD, Ph.D., of St. Boniface Hospital in Winnipeg, Canada, who was not involved with this research. "This study further highlights the need for the health care team to undertake a more detailed assessment of patients' physical fitness before the operation. The 6MWD is an important component of this evaluation."

According to Dr. Arora, the identification of patients at risk for POCD and other cognitive disorders should alert the health care team to consider modifying anesthetic and medication choices during-and-after the operations, as well as assist with discharge planning as patients transition to home. In addition, the health care team should consider strategies, such as prehabilitation, to optimize the patients' fitness before their operations. Dr. Arora explained that prehabilitation includes a combination of exercise training, education, and social support intended to improve patients' physical and psychological readiness for surgery.

"Prehabilitation may be of benefit to patients with poor physical fitness by improving postoperative recovery and post-discharge functional survival," said Dr. Arora. "Patient self-management and follow-through are essential, however, as is the patient's understanding of their health issues and their proposed plan of care."

Dr. Hayashi agrees that a multidisciplinary approach, which includes

elements such as prehabilitation, is key to a better assessment and treatment outcome. "Precise preoperative risk assessment for postoperative complications is critical, and when indicated, supervised exercise before an operation should be recommended to improve functional exercise capacity before heart surgery," he said.

More information: Kazuhiro Hayashi et al, Preoperative 6-Minute Walk Distance Is Associated With Postoperative Cognitive Dysfunction, *The Annals of Thoracic Surgery* (2018). [DOI: 10.1016/j.athoracsur.2018.03.010](https://doi.org/10.1016/j.athoracsur.2018.03.010)

Provided by Elsevier

Citation: Simple walking test helps predict risk for cognitive issues after heart surgery (2018, May 10) retrieved 19 April 2024 from <https://medicalxpress.com/news/2018-05-simple-cognitive-issues-heart-surgery.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.