

Study examines treatment options for women with recurrent ovarian cancer

May 23 2018



Intermediate magnification micrograph of a low malignant potential (LMP) mucinous ovarian tumour. H&E stain. The micrograph shows: Simple mucinous epithelium (right) and mucinous epithelium that pseudo-stratifies (left - diagnostic of a LMP tumour). Epithelium in a frond-like architecture is seen at the top of image. Credit: Nephron /Wikipedia. CC BY-SA 3.0

New research indicates that for women with advanced epithelial ovarian cancer whose cancer has relapsed after surgery, a second surgery is worth considering. The population-based registry study confirms the

results suggested by a recent randomized trial.

In the *Acta Obstetricia et Gynecologica Scandinavica* study, secondary [surgery](#) delayed a new recurrence for 2 years and resulted in at least 6 years median overall survival.

"Treatment for ovarian cancer is centralized to only a few hospitals in Norway. The favourable results may encourage other countries to adopt this model in order to treat more patients with secondary surgery at first recurrence," said lead author Dr. Witold Szczesny, of the Cancer Registry of Norway.

More information: Witold Szczesny et al, Survival after secondary cytoreductive surgery and chemotherapy compared with chemotherapy alone for first recurrence in patients with platinum-sensitive epithelial ovarian cancer and no residuals after primary treatment. A registry-based study, *Acta Obstetricia et Gynecologica Scandinavica* (2018). [DOI: 10.1111/aogs.13361](#)

Provided by Wiley

Citation: Study examines treatment options for women with recurrent ovarian cancer (2018, May 23) retrieved 18 April 2024 from <https://medicalxpress.com/news/2018-05-treatment-options-women-recurrent-ovarian.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.