


New approach to immunotherapy leads to complete response in breast cancer patient

June 4 2018



Three-dimensional culture of human breast cancer cells, with DNA stained blue and a protein in the cell surface membrane stained green. Image created in 2014 by Tom Misteli, Ph.D., and Karen Meaburn, Ph.D. at the NIH IRP.

A novel approach to immunotherapy developed by researchers at the National Cancer Institute (NCI) has led to the complete regression of breast cancer in a patient who was unresponsive to all other treatments. This patient received the treatment in a clinical trial led by Steven A. Rosenberg, M.D., Ph.D., chief of the Surgery Branch at NCI's Center for Cancer Research (CCR), and the findings were published June 4, 2018 in *Nature Medicine*. NCI is part of the National Institutes of Health.

"We've developed a high-throughput method to identify [mutations](#) present in a cancer that are recognized by the immune system," Dr. Rosenberg said. "This research is experimental right now. But because this new approach to immunotherapy is dependent on mutations, not on cancer type, it is in a sense a blueprint we can use for the [treatment](#) of many types of cancer."

The new immunotherapy approach is a modified form of adoptive cell transfer (ACT). ACT has been effective in treating melanoma, which has high levels of somatic, or acquired, mutations. However, it has been less effective with some common epithelial cancers, or cancers that start in the lining of organs, that have lower levels of mutations, such as stomach, esophageal, ovarian, and breast cancers.

In an ongoing phase 2 clinical trial, the investigators are developing a form of ACT that uses [tumor](#)-infiltrating lymphocytes (TILs) that specifically target tumor cell mutations to see if they can shrink tumors in patients with these common epithelial cancers. As with other forms of ACT, the selected TILs are grown to large numbers in the laboratory and are then infused back into the patient (who has in the meantime undergone treatment to deplete remaining lymphocytes) to create a stronger immune response against the tumor.

A patient with [metastatic breast cancer](#) came to the trial after receiving multiple treatments, including several chemotherapy and hormonal

treatments, that had not stopped her cancer from progressing. To treat her, the researchers sequenced DNA and RNA from one of her tumors, as well as normal tissue to see which mutations were unique to her cancer, and identified 62 different mutations in her tumor cells.

The researchers then tested different TILs from the patient to find those that recognized one or more of these mutated proteins. TILs recognized four of the mutant proteins, and the TILs then were expanded and infused back into the patient. She was also given the checkpoint inhibitor pembrolizumab to prevent the possible inactivation of the infused T cells by factors in the tumor microenvironment. After the treatment, all of this patient's cancer disappeared and has not returned more than 22 months later.

"This is an illustrative case report that highlights, once again, the power of immunotherapy," said Tom Misteli, Ph.D., director of CCR at NCI. "If confirmed in a larger study, it promises to further extend the reach of this T-cell therapy to a broader spectrum of cancers."

Investigators have seen similar results using mutation-targeted TIL treatment for patients in the same trial with other epithelial cancers, including liver cancer and colorectal cancer. Dr. Rosenberg explained that results like this in [patients](#) with solid epithelial tumors are important because ACT has not been as successful with these kinds of cancers as with other types that have more mutations.

He said the "big picture" here is this kind of treatment is not cancer-type specific. "All cancers have mutations, and that's what we're attacking with this immunotherapy," he said. "It is ironic that the very mutations that cause the cancer may prove to be the best targets to treat the [cancer](#)."

More information: Nikolaos Zacharakis et al, Immune recognition of

somatic mutations leading to complete durable regression in metastatic breast cancer, *Nature Medicine* (2018). [DOI: 10.1038/s41591-018-0040-8](#)

Provided by National Cancer Institute

Citation: New approach to immunotherapy leads to complete response in breast cancer patient (2018, June 4) retrieved 18 April 2024 from <https://medicalxpress.com/news/2018-06-approach-immunotherapy-response-breast-cancer.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.