


Asymmetrical neuron loss in Alzheimer's

June 29 2018



Credit: Christian Wachinger/LMU

As Alzheimer's disease progresses, neuron loss leads to marked lateral asymmetries in brain structure. A new LMU study has now identified genetic variants that promote the emergence of asymmetrical brain atrophy in Alzheimer's patients.

Alzheimer's disease leads to the loss of brain tissue, primarily in the hippocampus, which plays a crucial role in the formation of long-term memories. In previous studies, Christian Wachinger, a neuroinformatics specialist at LMU, was able to show that, as dementia progresses, asymmetries develop in the brain anatomy. These in turn reflect the asymmetrical loss of neurons. In a long-term study, he and his colleagues have tracked these in the shape of brain structures, and searched for genetic markers that might be functionally implicated in their emergence. As they report in the journal *Biological Psychiatry*, they have

indeed identified several statistically significant associations between specific genetic variants and progressive asymmetrical neuron loss in individuals with Alzheimer's disease.

Based at LMU's Clinic for Child and Adolescent Psychiatry, Psychosomatics and Psychotherapy, Wachinger leads a junior research group on artificial intelligence in medical imaging (AI-Med). The new study made use of 6395 magnetic resonance images acquired over a period of 9 years from 1241 [patients](#) who had been diagnosed with Alzheimer's disease. Using the program BrainPrint, which Wachinger developed earlier, the team quantified the degree of asymmetry of specific brain structures in the patient population at various timepoints. The anatomical results were then compared with genetic data from the same patients, focusing on single-site variants in 21 specific genes which had previously been linked with an increased risk for late-onset Alzheimer's disease in genome-wide association studies, and 10 others that appear to affect the relative volumes of certain brain substructures but had not been directly related to Alzheimer's risk per se. "The question we wanted to answer was whether there are genetic factors that make the brains of Alzheimer's patients particularly prone to asymmetrical neuron loss," Wachinger explains.

Wachinger's team indeed identified five genetic variants that are associated with the development of anatomical asymmetries in specific brain structures as Alzheimer's dementia progresses, while one other genetic polymorphism was found to promote asymmetry irrespective of [disease](#) status. "Our results provide new insights into the biological basis for the emergence of asymmetries in patients with Alzheimer's. They also yield new clues regarding the effects of known genetic risk factors on the anatomy of the [brain](#)," says Wachinger.

More information: Christian Wachinger et al. A Longitudinal Imaging Genetics Study of Neuroanatomical Asymmetry in Alzheimer's

Disease, *Biological Psychiatry* (2018). [DOI: 10.1016/j.biopsych.2018.04.017](https://doi.org/10.1016/j.biopsych.2018.04.017)

Provided by Ludwig Maximilian University of Munich

Citation: Asymmetrical neuron loss in Alzheimer's (2018, June 29) retrieved 3 May 2024 from <https://medicalxpress.com/news/2018-06-asymmetrical-neuron-loss-alzheimer.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--