

Bone mass may suffer when teenage girls binge drink

June 14 2018

Teenage girls who regularly binge drink may fail to reach their peak bone mass, according to a new study published in the *Journal of Studies on Alcohol and Drugs*.

The study, of 87 college [women](#), found that those who regularly binge drank in high school had lower [bone mass](#) in the spine. That was true even when researchers accounted for other factors that affect bone density—such as exercise, nutrition and smoking habits.

The findings suggest that poorer bone health can be added to the list of binge [drinking](#) risks for [young women](#), said lead researcher Joseph LaBrie, Ph.D., a professor of psychology at Loyola Marymount University, in Los Angeles, who conducted the research with health and human sciences professor Hawley Almstedt, Ph.D., R.D.N.

There are well-known short-term risks, LaBrie pointed out—such as alcohol poisoning, car accidents, poor academic performance and sexual assault.

"This study identifies a potential lifetime consequence of binge drinking in young women," he said.

The findings are based on female college students ages 18 to 20—a time when, LaBrie said, bone mass should still be accruing. Women generally reach their peak bone density at the spine between the ages of 20 and 25.

The study participants answered questionnaires about certain lifestyle factors and underwent measurements of their bone density in the lumbar spine. When it came to alcohol, the women were asked to think back to high school and report how often they'd binged—having four or more drinks within two hours.

Overall, LaBrie's team found, women who'd binged frequently since [high school](#) had lower bone mass than their peers. "Frequent" meant they'd binged at least 115 times—or nearly twice a month, on average.

The findings expand previous research linking heavy drinking to lower bone mass and higher fracture risk in older adults, suggesting that later in life bone issues may be linked to drinking early in life. Meanwhile, previous animal research has suggested that alcohol hinders healthy development of young bones.

LaBrie noted that anything that keeps a young woman from reaching her peak bone mass will probably raise her odds of developing osteoporosis years down the road.

For now, the findings offer girls and young women one more reason to avoid binge drinking and offers parents further support for seeking to delay onset of children's drinking.

"When we consider [bone](#) health," LaBrie said, "we always talk about things like exercise, calcium and vitamin D, and not smoking. We may also need to talk about avoiding [binge drinking](#)."

More information: Joseph W. Labrie et al, Heavy Episodic Drinking Is Associated With Poorer Bone Health in Adolescent and Young Adult Women, *Journal of Studies on Alcohol and Drugs* (2018). [DOI: 10.15288/jsad.2018.79.391](#)

Provided by Journal of Studies on Alcohol and Drugs

Citation: Bone mass may suffer when teenage girls binge drink (2018, June 14) retrieved 4 May 2024 from <https://medicalxpress.com/news/2018-06-bone-mass-teenage-girls-binge.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.