

Children with neuroblastoma have an elevated risk of long-term psychological difficulties

June 11 2018

A new study reveals that pediatric neuroblastoma patients are at elevated risk for long-term psychological impairment. In addition, those who experience such impairment as they get older tend to require special education services and to not go on to college. The findings are published early online in *Cancer*, a peer-reviewed journal of the American Cancer Society.

In the United States, neuroblastoma—a childhood <u>cancer</u> of nerve cells—is diagnosed at a <u>median age</u> of 17.3 months. Treatment advances in recent years have prolonged survival for many affected children, but their young age at diagnosis and the specific therapies they receive can make them vulnerable to health problems as their central nervous system develops.

To assess the long-term psychological effects of neuroblastoma and its treatment, Nina Kadan-Lottick, MD, MSPH, of Yale University School of Medicine and her colleagues studied 859 children who had been diagnosed with neuroblastoma at least five years earlier and were under 18 years old. Their median age at diagnosis was 0.8 years, and they were followed for a median of 13.3 years. These 859 neuroblastoma survivors were compared with 872 siblings of childhood cancer survivors.

Compared with siblings, neuroblastoma survivors had an increased prevalence of impairment in the domains of anxiety/depression (19



percent versus 14 percent), headstrong behavior (19 percent versus 13 percent), attention deficits (21 percent versus 13 percent), peer conflict/social withdrawal (26 percent versus 17 percent), and antisocial behavior (16 percent versus 12 percent).

Common treatments—vincristine, cisplatin, and retinoic acid—were not associated with impairment, but survivors who developed chronic health conditions as a result of their cancer treatment were at higher risk for developing worse outcomes. Specifically, developing pulmonary disease was linked with an increased risk of impairment in all five domains, and developing endocrine disease and peripheral neuropathy were each linked with impairment in three domains. In addition, survivors who experienced psychological impairment tended to require special education services and to not go on to college.

"These findings are novel because this is the first large study that could look at how <u>neuroblastoma</u> patients are doing in terms of psychological and educational outcomes. Before recent advances in treatment, this survivor population was much smaller and we were not able to analyze these sorts of long-term outcomes," said Dr. Kadan-Lottick. "The goal is not simply to get our patients to be cancer-free but also to optimize their mental, emotional, and social functioning as they move into adolescence and adulthood. Our hope is that these findings will help inform strategies for early screening and intervention to identify those survivors at highest risk for developing psychological and educational impairment later on in life."

June is National Cancer Survivors month.

More information: "Long-term psychological and educational outcomes in survivors of neuroblastoma: A report from the Childhood Cancer Survivor Study. Daniel J. Zheng, Kevin R. Krull, Yan Chen, Lisa Diller, Yutaka Yasui, Wendy Leisenring, Pim Brouwers, Rebecca



Howell, Jin-Shei Lai, Lyn Balsamo, Kevin C. Oeffinger, Leslie L. Robison, Gregory T. Armstrong, and Nina S. Kadan-Lottick. *Cancer*; Published Online: June 11, 2018, DOI: 10.1002/cncr.31379

Provided by Wiley

Citation: Children with neuroblastoma have an elevated risk of long-term psychological difficulties (2018, June 11) retrieved 3 May 2024 from https://medicalxpress.com/news/2018-06-children-neuroblastoma-elevated-long-term-psychological.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.