

Clinical outcomes and patient experiences vastly improved with hospital at home

June 25 2018

A new study to be published online June 25 in *JAMA Internal Medicine* reports that hospital at home (HaH) care provides a shorter length of stay; reductions in hospital readmissions, emergency department visits, and transfers to skilled nursing facilities; and, improved patient experience versus traditional inpatient care. The study, which spans nearly three years, includes patients with the broadest set of admitting diagnoses ever to be researched, thus strengthening the evidence base for hospital at home care.

Hospital at home provides acute [hospital](#)-level care in a patient's home as a substitute for traditional inpatient care. While HaH has been adopted in integrated [health care](#) systems and the Veterans Affairs health system, it has not been disseminated broadly due to the absence of a payment mechanism in fee-for-service Medicare. The authors note recent communication with the Secretary of the Department of Health and Human Services and progress towards the development of a payment model that the Centers for Medicare and Medicaid Services (CMS) will organize.

Study Overview

Researchers at the Icahn School of Medicine at Mount Sinai studied patients from November 2014 through August 2017. The study, funded by a Health Care Innovation Award from the Center for Medicare and Medicaid Innovation (CMMI) and The John A. Hartford Foundation,

compared outcomes of patients who received HaH care followed by a 30-day post-acute period of home-based transitional care with a control group of hospital inpatients who were HaH-eligible but refused participation or were seen in an [emergency department](#) when an HaH admission could not be initiated. Seeking to scale the hospital at home program, Mount Sinai entered into a partnership with Contessa Health, a health care company that manages acute-care services at home through bundled payment arrangements, in August 2017.

Results

The experiences of 507 patients were analyzed in the study, including 295 HaH patients and 212 controls. HaH participants, in contrast to controls, experienced the following outcomes:

- Shorter length of stay (3.2 days vs 5.5 days)
- Lower rates of hospital readmissions (8.6 percent vs. 15.6 percent)
- Lower rates of emergency department visits (5.8 percent vs. 11.7 percent)
- Fewer transfers to skilled nursing facilities (1.7 percent vs. 10.4 percent)
- More likely to rate their medical care highly (67.8 percent vs. 45.6 percent)
- More likely to rate their communications with nurses and doctors, in addition to communications about medicines, highly

"When you compare the patient treated in a hospital versus the home, the improvements are overwhelmingly positive from the point of view of clinical outcomes, patient safety, patient satisfaction, and cost savings," says Albert Siu, MD, MSPH, chair emeritus, Brookdale Department of Geriatrics and Palliative Medicine, Icahn School of Medicine at Mount Sinai. Broad dissemination of hospital at home, though, will require a

funding mechanism. "The availability of a payment mechanism is the one obstacle that stands in the way of widespread adoption of hospital at [home care](#)," says Dr. Siu.

On June 14, the Secretary of the Department of Health and Human Services (HHS) indicated that the HHS was "keenly interested in ideas for home-based, hospital-level care" and asked the Mount Sinai researchers to work with CMS as it develops a model for hospital care in the home. "We are pleased that the Secretary has asked CMS to explore a payment model that will leverage the work we've done over the past three years. A new payment model would add tremendously to Medicare's portfolio of shared savings programs," says Dr. Siu

Dr. Siu and his team estimate that 575,000 patients in the United States could qualify for HaH care every year. "If we treated just one in five of those patients, Medicare would save about \$45 million a year," says Dr. Siu. Reasons for greater efficiencies include shorter lengths of stay when patients are treated at home, reduced [hospital readmissions](#), lower rates of adverse events, and fewer emergency department revisits.

Treating patients at home is part of a much broader drive by Mount Sinai, a health system encompassing seven hospitals in Manhattan, to develop its hospitals into centers of care for the broader community. "The hospital of the future must closely mirror the needs of its community," says Kenneth L. Davis, MD, President and CEO of Mount Sinai. "Knowing when a patient will be better off receiving hospital-level care at home is key. Innovative care models, like hospital at home, ensure patients get the same level of care at home and avoid costly and unnecessary hospitalization."

How Does Hospital at Home Work?

Hospital at home care provides hospital-level services for beneficiaries

with selected acute illnesses who would otherwise be hospitalized. Typically, a patient is admitted while in the hospital, then sent home via ambulance, where a physician or nurse practitioner provides home-based acute care services. Nurses visit once or more a day to provide most of the care and a physician or NP sees [patients](#) at least daily in person or via video call facilitated by the nurse. Durable medical equipment, phlebotomy, and home radiography are provided when needed. Once the patient has recovered from the acute illness, he or she is discharged and the 30-day post-acute period begins. During this period, nurses and social workers provide self-management support and coordination of care with primary care clinicians, specialists, rehabilitation, and outpatient testing as needed.

Provided by The Mount Sinai Hospital

Citation: Clinical outcomes and patient experiences vastly improved with hospital at home (2018, June 25) retrieved 4 May 2024 from <https://medicalxpress.com/news/2018-06-clinical-outcomes-patient-vastly-hospital.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--