

Common diabetes drug found safe for most diabetics with kidney disease

June 5 2018



Oral diabetes drug metformin is safe for most diabetics who also have chronic kidney disease (CKD). Credit: Johns Hopkins Medicine

Results of a large-scale study suggest that the oral diabetes drug metformin is safe for most diabetics who also have chronic kidney



disease (CKD). The study of more than 150,000 adults by Johns Hopkins Medicine investigators found that metformin's association with the development of a life-threatening condition called lactic acidosis was seen only among patients with severely decreased kidney function.

A report on the study appeared online June 4 in JAMA Internal Medicine.

Metformin is the first-line medication for treatment of type 2 <u>diabetes</u> in adults and the fifth most commonly used drug in the United States. However, there has been a longstanding concern in the medical community about prescribing metformin for people with both diabetes and CKD because it may cause <u>lactic acidosis</u>, says senior study author Morgan Grams, M.D.,Ph.D., M.H.S., an associate professor of medicine and epidemiology at the Johns Hopkins University School of Medicine. Lactic acidosis is a rare but serious complication, occurring when lactate—a product of glucose breakdown—builds up in the bloodstream, producing symptoms such as severe muscle pain, cramps, nausea and weakness. Because <u>patients</u> with CKD may be at increased risk of metformin-associated lactic acidosis, health care providers historically avoided prescribing metformin in this population.

The Food and Drug Administration recently revised its labeling of metformin so the drug could be used more by CKD patients, says Grams, and some regulatory and professional society guidelines cautiously support use of the drug by diabetic patients with moderate kidney disease. But data addressing the drug's safety for patients with moderate to severe kidney disease had been inconclusive, she says.

A better assessment of acidosis risk could "provide reassurance that millions of people with diabetes and CKD may be able to safely use metformin," Grams says. Past studies show that metformin has multiple benefits beyond controlling blood sugar, including less weight gain, a lower risk of heart attack and even higher long-term survival than other



medications that patients take to control blood sugar. About 380 million people worldwide and 29 million people in the United States are affected by diabetes. Approximately 19 percent—or around 5 million—of those in the U.S. with diabetes also have CKD, Grams says.

"Our study demonstrates that the first-line and common diabetes medication is safer in patients with CKD than once thought," Grams says. "From a public health perspective, the potential benefits of using metformin for patients with diabetes and CKD are vast, given the increasing number of people affected with both diseases worldwide."

For the new study, Grams and her co-authors analyzed 2004-2017 medical records from a group of 75,413 patients with diabetes managed by Geisinger Health System in Pennsylvania. They looked at patients' estimated glomerular filtration rates (eGFRs—a measure of <u>kidney</u> <u>function</u>), medications prescribed for diabetes management, hospitalizations for acidosis, demographic information, cigarette smoking status and presence of cardiovascular disease.

The research team used computer-based statistical models to analyze the risk of developing acidosis among metformin users compared with nonusers. The researchers adjusted for conditions such as cardiovascular disease and smoking status.

Overall, Grams and her colleagues found an association between metformin use and acidosis only in patients with severely decreased kidney function, defined as an eGFR of less than 30 mL/min/1.73 m2. A normal eGFR is considered to be over 90 mL/min/1.73 m2. The researchers observed more than double the risk in patients with severely decreased kidney function who used another type of diabetes medication.

In addition, the researchers compared new metformin users with new



users of another class of diabetes medications called sulfonylureas within different categories of CKD stages. They replicated their results in a group of 82,017 patients from MarketScan—a database of inpatient and outpatient claims data from 350 private health systems—studying records from 2010 to 2014.

The Geisinger patients averaged age 60, and 51 percent were female. Of the patients, 45 percent (34,095) were taking metformin at enrollment in the study and 13,781 of the remaining patients were subsequently prescribed metformin during a 5.7-year follow-up period.

Among the Geisinger participants, there were 2,335 hospitalizations for acidosis over the study period. Compared with alternative forms of blood sugar management, metformin use was not associated with acidosis overall, nor was it associated with acidosis among patients with eGFR levels of 45 to 59 mL/min/1.73 m2 (considered mildly to moderately decreased kidney function) or among patients with eGFR levels of 30 to 44 mL/min/1.73 m2 (considered moderately to severely decreased kidney function).

However, metformin use was associated with an increased risk of acidosis at an eGFR level of less than 30 mL/min/1.73 m2.

These results were consistent when new metformin users were compared with new sulfonylurea users and with the MarketScan group of patients. There were 67,578 new metformin users in this group, and 14,439 new sulfonylurea users. There were 238 acidosis events among metformin users and 94 among sulfonylurea users. Lower eGFR was a risk factor for acidosis in both groups.

"Our results support cautious use of metformin in patients with type 2 diabetes and eGFR of at least 30 mL/min/1.73 m2," Grams says.



While the researchers say these data are promising, they caution that patients receiving metformin could have different underlying acidosis risk than patients who did not receive <u>metformin</u>, and that most of the study sample was white, which may limit generalizability by race.

Metformin, beyond having potential health benefits, is relatively inexpensive. The medication can be obtained for \$4 per month for the 500 milligram dose. In contrast, some of the newer diabetes agents can cost more than \$400 per month.

More information: Benjamin Lazarus et al, Association of Metformin Use With Risk of Lactic Acidosis Across the Range of Kidney Function, *JAMA Internal Medicine* (2018). DOI: <u>10.1001/jamainternmed.2018.0292</u>

Provided by Johns Hopkins University School of Medicine

Citation: Common diabetes drug found safe for most diabetics with kidney disease (2018, June 5) retrieved 28 April 2024 from https://medicalxpress.com/news/2018-06-common-diabetes-drug-safe-diabetics.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.