

No evidence that vitamin D protects against high blood pressure in pregnancy

June 20 2018



Credit: CC0 Public Domain

There is no strong evidence that vitamin D protects against pregnancy-induced high blood pressure (hypertension) or pre-eclampsia, conclude researchers in *The BMJ* today.

The findings support current World Health Organization guidance that evidence recommending vitamin D supplements for [women](#) during pregnancy to reduce adverse pregnancy outcomes is insufficient. However, in many countries, including the UK and the US, [pregnant women](#) are advised to take a daily dose of vitamin D.

It is common for pregnant women to have low levels of vitamin D, which can suppress the hormone that regulates [blood pressure](#) which may increase the risk of both [hypertension](#) and pre-eclampsia during pregnancy.

Previous population-based studies have found that women with lower levels of vitamin D are at greater risk of pre-eclampsia and some trials of vitamin D supplementation in pregnancy suggest a potential benefit. But it remains unclear whether vitamin D is a cause of pre-eclampsia.

So an international team led by Maria Magnus at the University of Bristol, set out to investigate whether vitamin D has an effect on [pregnancy induced hypertension](#) or pre-eclampsia.

Using a technique called Mendelian randomisation, they examined whether genetic variants associated with vitamin D production and metabolism also influenced the risk of pregnancy-induced hypertension and pre-eclampsia for 7,389 women (751 with gestational hypertension and 135 with pre-eclampsia) from two large European studies (Avon Longitudinal Study of Parents and Children, also known as Children of the 90s, and Generation R Study).

They also performed another (two sample) Mendelian randomisation analysis of 3,388 pre-eclampsia cases and 6,059 controls.

Analysing genetic information as proxies for the exposure of interest in this way avoids some of the problems that afflict traditional

observational studies, making the results less prone to unmeasured (confounding) factors. An association that is observed using Mendelian randomisation therefore strengthens the inference of a causal relationship.

Mendelian randomisation analysis showed no evidence to support a direct (causal) effect of vitamin D levels on risk of gestational hypertension or pre-eclampsia.

The researchers point to some study limitations. For example, the analyses were restricted to pregnant women. If vitamin D affects fertility, this may have resulted in selection bias.

In light of the uncertainty, they suggest further studies with a larger number of women with preeclampsia or more genetic variants that would increase the predictive power of vitamin D levels are needed.

"In combination with adequately powered clinical trials, this could help finally establish whether [vitamin D](#) status has a role in [pregnancy](#) related hypertensive disorders." they conclude.

More information: Vitamin D and risk of pregnancy related hypertensive disorders: mendelian randomisation study, www.bmj.com/content/361/bmj.k2167

Provided by British Medical Journal

Citation: No evidence that vitamin D protects against high blood pressure in pregnancy (2018, June 20) retrieved 25 April 2024 from <https://medicalxpress.com/news/2018-06-evidence-vitamin-d-high-blood.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.