


# Functional MRI reveals memory in sleeping toddlers

June 4 2018

---



It is difficult to study brain activity in very young children. Researchers at the UC Davis Center for Mind and Brain have come up with a protocol to use functional MRI scanning with children under two years old while they are asleep. This fMRI image shows activity in the hippocampus, a region of the brain associated with memory formation and recall. Credit: Simona Ghetti, UC Davis

Our ability to remember past events develops rapidly in the first couple

of years of life, but it's not clear exactly how this happens. Researchers at the Center for Mind and Brain at the University of California, Davis have now been able to carry out functional MRI brain scans of sleeping toddlers, and show for the first time how specific brain regions are activated during memory recall in two-year-olds.

The team is led by Professor Simona Ghetti, Department of Psychology and Center for Mind and Brain. Ghetti's laboratory is interested in how [memory](#) and cognition develop in children.

The hippocampus, a small structure deep in the middle of the brain, is important for forming and recalling memories and is thought to be involved in developing memory abilities in infants.

Functional Magnetic Resonance Imaging (MRI) is a powerful technique for measuring activity in the brain. But it's difficult to conduct conventional fMRI studies with very young children, because it requires completing a task while lying still with your head inside a large, noisy scanning machine.

Ghetti and postdoctoral researcher Janani Prabhakar worked out an alternative protocol that allows them to take measurements from a child during natural, [nighttime sleep](#) in the MRI scanner.

## **Asleep inside a scanner**

Children first listened to a [song](#) while playing with toys in the lab, so they formed memories of hearing the song while playing with a stuffed dog.

At their usual bedtime, they went to sleep inside the MRI scanner. While the child slept, the researchers played recordings of songs that they had heard earlier or unfamiliar songs.

The toddlers were also tested, while awake, on whether they could remember in which room they heard the song, or if they could associate the song with a toy they were playing with when they heard it for the first time. This ability to remember things in context—where it happened, what you were doing at the time—is called episodic memory.

Prabhakar and Ghetti found a stronger signal from the hippocampus when a familiar song was played. Children who showed good episodic memory also showed stronger hippocampal activation in the fMRI test, they found.

**More information:** Janani Prabhakar et al., "Memory-related hippocampal activation in the sleeping toddler," *PNAS* (2018).  
[www.pnas.org/cgi/doi/10.1073/pnas.1805572115](http://www.pnas.org/cgi/doi/10.1073/pnas.1805572115)

Provided by UC Davis

Citation: Functional MRI reveals memory in sleeping toddlers (2018, June 4) retrieved 19 April 2024 from <https://medicalxpress.com/news/2018-06-functional-mri-reveals-memory-toddlers.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.