

161 genetic factors for myopia identified

June 15 2018

The international Consortium for Refractive Error and Myopia (CREAM) recently published the largest-ever genetic study of myopia in *Nature Genetics*. Researchers from the Gutenberg Health Study at the Medical Center of Johannes Gutenberg University Mainz participated in this study, which identified 161 genetic factors for myopia. This quadruples the number of known genetic risk factors playing a role in all retinal cell types. Most of them are involved in processing light. This supports the assumption that insufficient sunlight is an important trigger in the development of myopia.

Myopia, also known as short-sightedness or nearsightedness, is the most common disorder affecting the eyesight, and it is on the increase. This is particularly worrying for severely short-sighted people as it increases their risk of developing eyesight complications. The causes are both genetic and environmental.

The international research group CREAM, which includes scientists from the Gutenberg Health Study of the Mainz University Medical Center, has now made important progress toward understanding the mechanisms behind the development of the condition. They evaluated the data of more than 250,000 people from Europe, Asia, and North America in cooperation with gene test provider 23andMe.

The study established 161 [genetic factors](#) for spherical equivalent and [myopia](#), most of which were previously unknown. It became clear that all retinal cell types play a role in the development of myopia alongside their primary role as light processors. This supports the theory that the

internal layer of the eye communicates with the external layer to increase the length of the eye, which is a decisive factor in the development of myopia.

"We have known for some time that education-related behavior is a major environmental factor in developing short-sightedness," said co-author Professor Norbert Pfeiffer, head of the Department of Ophthalmology at the Mainz University Medical Center. It is unclear whether close-up reading or a lack of sunlight plays a larger role. The new results provide important insights into the underlying biological mechanisms. They also support the most important advice Pfeiffer can give to concerned parents as a preventive measure against myopia: "Send your kids to play outside for two hours every day. And it's not just their eyes that will benefit."

The spread of short-sightedness is a worldwide phenomenon, particularly in Southeast Asia, where the incidence of myopia in school children has increased notably over the last decades. This is likely due to rising levels of education. People who read a great deal also perform a lot of close-up work, usually in poor levels of daylight. The eye adjusts to these visual habits and the eyeball becomes more elongated than normal as a result. But if it becomes too elongated, the cornea and lens focus the image just in front of the retina instead of on it, making distant objects appear blurry.

More information: Genome-wide association meta-analysis highlights light-induced signaling as a driver for refractive error, *Nature Genetics* (2018). [DOI: 10.1038/s41588-018-0127-7](https://doi.org/10.1038/s41588-018-0127-7)

Provided by Universitaet Mainz

Citation: 161 genetic factors for myopia identified (2018, June 15) retrieved 9 April 2024 from <https://medicalxpress.com/news/2018-06-genetic-factors-myopia.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.