

## Infant omega-3 supplementation tied to decreased waist size

June 8 2018

---



(HealthDay)—Omega-3 long-chain polyunsaturated fatty acids (n-3

LCPUFA) supplementation in infancy is associated with reduced insulin concentrations and insulin resistance in boys and reduced waist circumference in both sexes at age 5 years, according to a study published online June 8 in *Pediatrics*.

Valene H.L. See, Ph.D., from the Royal Perth Hospital in Australia, and colleagues randomly assigned 420 infants to a daily supplement of n-3 LCPUFA or olive oil (control) from birth to 6 months. Growth, body composition, and cardiometabolic risk factors were evaluated at 5 years of age.

The researchers found that infants who received n-3 LCPUFA had a smaller waist circumference at 5 years (coefficient, 1.1 cm), which remained significant after adjusting for confounders (coefficient, 0.8 cm). Boys who received n-3 LCPUFA supplementation had a 21 percent reduction in [insulin](#) concentrations and a 22 percent reduction in [insulin resistance](#) versus the control group. At birth, 2.5 years, and 5 years, there were no other differences in growth and cardiometabolic risk factors between the groups.

"Longer-term follow-up of the cohort is warranted to determine whether these differences are maintained into adolescence," write the authors.

**More information:** [Abstract/Full Text \(subscription or payment may be required\)](#)

Copyright © 2018 [HealthDay](#). All rights reserved.

Citation: Infant omega-3 supplementation tied to decreased waist size (2018, June 8) retrieved 25 April 2024 from <https://medicalxpress.com/news/2018-06-infant-omega-supplementation-tied-decreased.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.