

Optimal sleep linked to lower risks for dementia and early death

June 6 2018



Credit: CC0 Public Domain

Short and long daily sleep duration were risk factors for dementia and premature death in a study of Japanese adults aged 60 years and older. The findings are published in the *Journal of the American Geriatrics*

Society.

Among 1,517 adults who were followed for 10 years, 294 developed [dementia](#) and 282 died. Age- and sex-adjusted incidence rates of dementia and all-cause mortality were greater in those with daily sleep duration of less than 5.0 hours and 10.0 hours or more, compared with those with daily sleep duration of 5.0 to 6.9 hours. Participants with short sleep duration who had high physical activity did not have a greater risk of dementia and death, however.

"Given the beneficial effects of [physical activity](#) on risk of sleep disturbance, these findings indicate that not only maintenance of appropriate sleep duration, but also modification of lifestyle behaviors related to sleep may be an effective strategy for preventing dementia and [premature death](#) in elderly adults," the authors wrote.

More information: *Journal of the American Geriatrics Society* (2018). DOI: [10.1111/jgs.15446](https://doi.org/10.1111/jgs.15446) , onlinelibrary.wiley.com/journal/15325415

Provided by Wiley

Citation: Optimal sleep linked to lower risks for dementia and early death (2018, June 6) retrieved 9 April 2024 from <https://medicalxpress.com/news/2018-06-optimal-linked-dementia-early-death.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--