


First photoactive drug to fight Parkinson's disease

June 8 2018



Credit: Universidad de Barcelona

An international team has designed the first potentially therapeutic photoactive drug, MRS7145, to fight Parkinson's disease, according to the new article in *Journal of Controlled Release*.

This compound, which proved effective in laboratory animals in vivo, has been carried out by a team led by Francisco Ciruela, from the Faculty of Medicine and Health Sciences of the University of Barcelona, and collaborators.

Optopharmacology: beyond the limits of conventional

drugs

Parkinson's is the second most common neurodegenerative disease after Alzheimer's and it affects more than 1 percent of the population. This disease, which affects the central nervous system, affects more than six million people worldwide, figures that could go higher than twelve million by 2030, according to data from the World Health Organization (WHO). In this disease, the neurotransmitter that controls the motor activity, dopamine, is reduced due the progressive death of dopaminergic neurons.

The action of [conventional drugs](#) is sometimes limited due several factors –lack of spatial specificity, slow and inaccurate distribution, etc.- which can reduce its therapeutic efficiency. Also, the efficiency of the traditional treatment on Parkinson's (levodopa) diminishes over time and it requires to increase the administration of the dose or to change the drug. The [adverse effects](#) of new drugs (uncontrolled movements in the body, motor fluctuations, etc.) are another common denominator in most patients.

Optopharmacology is an innovative discipline based on the use of light –with a certain wave length- to control the activity of drugs. Therefore, light-sensitive drugs can act with a higher spatial and time precision and without creating adverse effects in the body.

MRS7145: opening the way to treat neurodegenerative diseases

The MRS7145, the first potentially therapeutic photoactive drug to fight Parkinson's, is a photo-sensitive derivate from SCH442416, a selective antagonist of adenosine A2A receptor. In scientific bibliography, some A2A antagonist receptors came up as potential drugs to fight

Parkinson's, since they take part in the involved mechanisms in controlling the movement.

This photoactive drug is an inactive chemical compound that gets activated with light from a visible spectrum (with a 405 nm wave length) which is not harmful for the body. A series of optical fibers –planted in the striated bodies of laboratory animals- provide the irradiation of this area in the brain, which is responsible for the control of motor activity.

Lecturer Francisco Ciruela says, "once the striated body is radiated with violet light the active drug is released and blocks the adenosine A2A receptor. The blocking of adenosine receptors has an administrating effect on the activity of dopamine (pro-dopaminergic action)."

Photo-sensitive molecules that improve patients' quality of life

Improving the spatial and temporary precision of the drug and strengthening the commitment of the patient to the therapy are some of the benefits of optopharmacology in Parkinson's. "A fine time-space precision will enable manipulating the neural circuits in detail and set the functioning of those with therapeutic and neuroprotective purposes," says Ciruela.

"Nowadays, in addition, there are treatments that are based on the implementation of electrodes in the brain of patients with Parkinson's to control the electric activity of neurons. In the same lines, optical fibers could make light getting to almost any part of the body (spatial resolution), and these organs would be radiated with light controlled by an electronic device that would regulate the intensity and length of radiation (time resolution).

Maintaining the commitment of patients to the set therapeutic guideline on the long run is a big challenge for chronic diseases. "With a slow release system from the photoactive drug, such as a coupled patch with a radiation system remotely controlled by a phone App, the doctor could control in a precise manner the release of the most efficient dose of the active drug in the place of action," says Ciruela.

Although the clinical application of this photoactive [drug](#) in patients is still far, this pharmacological innovation could lead the way to a research on new therapeutical solutions for this chronic disease. The new article, published in *Journal of Controlled Release*, is therefore a step forward in the field of pharmacology to design new therapeutical strategies with photo-sensitive molecules and to set innovative clinical protocols to improve patients' quality of life.

More information: Jaume Taura et al. Remote control of movement disorders using a photoactive adenosine A_{2A} receptor antagonist, *Journal of Controlled Release* (2018). [DOI: 10.1016/j.jconrel.2018.05.033](#)

Provided by University of Barcelona

Citation: First photoactive drug to fight Parkinson's disease (2018, June 8) retrieved 25 April 2024 from <https://medicalxpress.com/news/2018-06-photoactive-drug-parkinson-disease.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--