

Physical fitness and healthy aorta key to brain ageing

June 12 2018



Credit: CC0 Public Domain

The rate of decline in certain aspects of memory may be explained by a combination of overall physical fitness and the stiffness of the central arteries, researchers from Swinburne's Centre for Human

Psychopharmacology have found.

A study to be published in the *Journal of Alzheimer's Disease* considers the mechanisms underlying [cognitive performance](#) in [older people](#) living independently.

Lead author, Ph.D. candidate Greg Kennedy, says that from early adulthood, memory and other aspects of cognition slowly decline, with an increasing risk of developing into dementia later in life.

"Exactly why this occurs is unclear, but research indicates that exercise and [physical fitness](#) are protective," Mr Kennedy says. "A healthier, more elastic aorta is also theorised to protect cognitive function, by reducing the negative effects of excessive blood pressure on the brain."

The study investigated whether fitness was associated with better cognition through a healthier aorta.

Physical fitness and arterial stiffness assessment

One hundred and two people (73 females and 29 males), aged between 60 and 90 years, living independently in aged care communities, were recruited in Melbourne, Australia.

Their fitness was assessed with the Six-Minute Walk test which involved participants walking back and forth between two markers placed 10 metres apart for six minutes.

Only participants who completed the full six minutes were included in the analysis, which assessed the [stiffness](#) of their arteries and cognitive performance.

The researchers found that (along with Body Mass Index and sex) the

combination of fitness and aortic stiffness explained a third of the variation in performance in working memory in older people.

"People generally are less fit and have stiffer arteries as they age, which seems to explain the difference in memory ability that is usually attributed to 'getting older'," Mr Kennedy says.

Interestingly, physical fitness did not seem to affect central arterial stiffness, however Mr Kennedy points out that only current fitness was assessed – long term [fitness](#) may be a better predictor of central arterial stiffness, however this has yet to be investigated.

"Unfortunately, there is currently no effective pharmacological intervention that has proven effective in the long term in reducing this decline or staving off dementia," Mr Kennedy says.

"The results of this study indicate that remaining as physically fit as possible, and monitoring central arterial health, may well be an important, cost effective way to maintain our [memory](#) and other brain functions in older age."

More information: Greg Kennedy et al, Physical Fitness and Aortic Stiffness Explain the Reduced Cognitive Performance Associated with Increasing Age in Older People, *Journal of Alzheimer's Disease* (2018). DOI: 10.3233/JAD-171107 , [dx.doi.org/10.3233/JAD-171107](https://doi.org/10.3233/JAD-171107)

Provided by Swinburne University of Technology

Citation: Physical fitness and healthy aorta key to brain ageing (2018, June 12) retrieved 10 April 2024 from <https://medicalxpress.com/news/2018-06-physical-healthy-aorta-key-brain.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.