

# Teens with concussion may benefit from earlier physical therapy

June 27 2018

---

For adolescents with symptoms following a concussion, starting physical therapy (PT) earlier—within less than three weeks after the injury—provides outcomes similar to those of later PT, suggests a study in the July issue of *The Journal of Neurologic Physical Therapy (JNPT)*.

"Multimodal PT interventions administered by licensed physical therapists may be feasible and safe even within the first few weeks after injury to help facilitate prompt recovery and mitigate the onset of secondary effects from delayed treatment," write Catherine Quatman-Yates, DPT, Ph.D., of The Ohio State University, Columbus, and colleagues. The study is part of a *JNPT* special issue on "Rehabilitation Management of Concussion," highlighting research-driven changes geared toward promoting return to activity in young patients with [concussion](#).

## Similar Outcomes for Teens with Concussion Undergoing Earlier or Later PT

The researchers looked how the timing of PT affected the course of concussion-related symptoms in 120 adolescents: 78 females and 42 males, median age 14 years. Physical therapy was classified as early (beginning 0 to 20 days after concussion) in 27.5 percent of patients, middle (21 to 41 days) in 32.5 percent, and late (42 days or after) in 40 percent.

The PT program consisted of progressive exercise; vestibular/oculomotor training (targeting inner ear/balance and visual symptoms); and cervical spine manual therapy, stretching, and strengthening exercises. This multimodal treatment was delivered by licensed [physical therapists](#) with special training in concussion treatment.

Whether started earlier or later, PT led to similar reductions in concussion-related symptoms. The number of sessions and duration of PT care were similar across groups. There was a low rate of adverse events, most of which were unrelated to PT.

Symptoms worsened in a few patients, more commonly in the late PT group. Some of these patients may have had concussion-related impairments not directly addressed by PT, such as anxiety, depression, or sleep problems.

Recent research has led to new insights into medical management of concussion in children and adolescents. Past guidelines recommended complete physical and cognitive (mental) rest after concussion, until symptoms resolved. But recent studies have suggested that resting for more than a day or two has limited benefits, and may even be linked to increased concussive symptoms.

Today, [concussion management](#) is shifting toward a shorter period of rest, followed by gradual return to usual activities, guided by the patient's symptoms. Physical therapy has been recommended for adolescents with persistent symptoms of concussion, generally after three weeks.

The new study provides evidence that starting PT earlier is a safe and feasible approach for adolescents after concussion, with improved symptoms regardless of the timing of the intervention. "Introducing PT earlier in the recovery process may be beneficial in minimizing the potential burden of longer recovery trajectories," Dr. Quatman-Yates

and coauthors write. They emphasize the need for further research to determine PT's role in the "optimal plan of care" for young patients with concussion.

Other articles in the special issue include a neuroscience perspective on the role of rest versus physical activity in recovery for young people with concussion, along with new research on changes in vestibular/oculomotor function and the role of balance testing after concussion.

Physical therapists can play a critical role in evaluating and choosing targeted interventions most likely to result in the best outcomes for patients with concussion, according to a Guest Editorial by Karen L. McCulloch, PT, Ph.D., NCS, and Kathleen Gill-Body, PT, DPT, MS, NCS, FAPTA. They write, "We are in an ideal position to continue our process of returning people to activities and roles that they care about... because it is what we do."

**More information:** Anne Lennon et al. An Exploration of the Impact of Initial Timing of Physical Therapy on Safety and Outcomes After Concussion in Adolescents, *Journal of Neurologic Physical Therapy* (2018). [DOI: 10.1097/NPT.0000000000000227](https://doi.org/10.1097/NPT.0000000000000227)

Provided by Wolters Kluwer Health

Citation: Teens with concussion may benefit from earlier physical therapy (2018, June 27) retrieved 24 May 2024 from <https://medicalxpress.com/news/2018-06-teens-concussion-benefit-earlier-physical.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
---