

## Tonsil and adenoid removal associated with respiratory, allergic and infectious disease

June 7 2018

Tonsil and adenoid removal associated with long-term risks of respiratory, allergic and infectious diseases Removing tonsils and adenoids in childhood increases the long-term risk of respiratory, allergic and infectious diseases, according to researchers who have examined—for the first time—the long-term effects of the operations.

The researchers suggest renewed evaluation of alternatives to these common paediatric surgeries that include removal of tonsils (tonsillectomy) to treat chronic tonsillitis or adenoids (adenoidectomy) to treat recurrent <u>middle ear</u> infections.

The adenoids and tonsils are strategically positioned in the nose and throat respectively to act as a first line of defense, helping to recognise airborne pathogens like bacteria and viruses, and begin the immune response to clear them from the body.

The collaborative study initiated by the Copenhagen Evolutionary Medicine program looked at the <u>long-term effects</u> of removing the tonsils and adenoids in childhood, compared with children who had not undergone the surgeries.

University of Melbourne researcher Dr. Sean Byars and Professor Jacobus Boomsma from the University of Copenhagen led the research, with Professor Stephen Stearns from Yale University. The research is published in the *Journal of the American Medical Association Otolaryngology Head and Neck Surgery*.



The team analysed a dataset from Denmark of 1,189,061 children born between 1979 and 1999, covering at least the first 10 years and up to 30 years of their life. Of the almost 1.2 million children, 17,460 had adenoidectomies, 11,830 tonsillectomy and 31, 377 had adenotonsillectomies, where both tonsils and adenoids removed. The children were otherwise healthy.

"We calculated <u>disease</u> risks depending on whether adenoids, tonsils or both were removed in the first 9 years of life because this is when these tissues are most active in the developing immune system," Dr. Byars said.

The analysis showed:

- Tonsillectomy was associated with an almost tripled relative risk—the risk for those who had the operation compared with those who didn't—for diseases of the <u>upper respiratory tract</u>. These included asthma, influenza, pneumonia and chronic obstructive pulmonary disorder or COPD, the umbrella term for diseases such as chronic bronchitis and emphysema.
- The absolute risk (which takes into account how common these diseases are in the community) was also substantially increased at 18.61 percent.
- Adenoidectomy was found to be linked with a more than doubled relative risk of COPD and a nearly doubled relative risk of upper respiratory tract diseases and conjunctivitis. The absolute risk was also almost doubled for upper respiratory diseases but corresponded to a small increase for COPD, as this is a rarer condition in the community generally.

"The association of tonsillectomy with respiratory disease later in life may therefore be considerable for those who have had the operation," Prof Boomsma said.



The team delved deeper into the statistics to reveal how many operations needed to be performed for a disease to occur at a greater rate than normal, known as the number needed to treat or NNT.

"For tonsillectomy, we found that only five people needed to have the operation to cause an extra upper respiratory disease to appear in one of those people," added Prof Boomsma.

The team also analysed conditions that these surgeries directly aimed to treat, and found mixed results:

- Adenoidectomy was associated with a significantly reduced risk for sleep disorders and all surgeries were associated with significantly reduced risk for tonsillitis and chronic tonsillitis, as these organs were now removed.
- However, there was no change in abnormal breathing up to the age of 30 for any <u>surgery</u> and no change in sinusitis after tonsillectomy or adenoidectomy.
- Following adenotonsillectomy the relative risk for those who had the operation was found to increase four or five-fold for otitis media (inflammation of the middle ear) and sinusitis also showed a significant increase.

The study suggests that shorter-term benefits of these surgeries may not continue up to the age of 30 apart from the reduced risk for tonsillitis (for all surgeries) and sleep disorders (for adenoidectomy).

Instead, the longer-term risks for abnormal breathing, sinusitis and otitis media were either significantly higher after surgery or not significantly different.

The researchers note that there will always be a need to remove tonsils and adenoids when those conditions are severe.



"But our observed results that show increased risks for long-term diseases after surgery support delaying tonsil and <u>adenoid</u> removal if possible, which could aid normal immune system development in childhood and reduce these possible later-life disease risks, Dr. Byars said.

"As we uncover more about the function of immune tissues and the lifelong consequences of their removal, especially during sensitive ages when the body is developing, this will hopefully help guide treatment decisions for parents and doctors."

**More information:** *Journal of the American Medical Association Otolaryngology Head and Neck Surgery*, <u>DOI:</u> <u>10.1001/jamaoto.2018.0614</u>

Provided by University of Melbourne

Citation: Tonsil and adenoid removal associated with respiratory, allergic and infectious disease (2018, June 7) retrieved 5 May 2024 from <u>https://medicalxpress.com/news/2018-06-tonsil-adenoid-respiratory-allergic-infectious.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.