

Almost half of US adults who drink, drink too much, and continue to do so

July 17 2018



Credit: CC0 Public Domain

A new study led by Boston University School of Public Health (BUSPH) researchers has found that about 40 percent of adults in the United States who drink alcohol do so in amounts that risk health consequences, and

identifies a range of factors associated with starting or stopping drinking too much.

The study, published in the *Journal of Substance Use*, found that 73 percent of those drinking risky amounts were still doing so two to four years later, while 15 percent of those not drinking risky amounts began to. Starting to drink too much was associated with being younger, transitioning to [legal drinking age](#), being male and white, and smoking and drug use, among other social factors.

"Some people just stop drinking too much, but most continue for years, and others not drinking too much will begin doing so during adulthood," says lead author Richard Saitz, professor of community health sciences at BUSPH. "Public health and clinical messages need repeating, particularly in [young adulthood](#). Once is not enough."

The researchers used data collected by interview from a nationally representative sample of more than 34,000 adults in the US who completed the National Epidemiologic Survey on Alcohol and Related Conditions in 2001-2002 and again in 2004-2005.

The survey assessed participants' drinking in the past month using a well-validated interview tool. "At-risk use" was defined as more than 14 drinks per week on average or more than 4 on an occasion for men, and more than 7 per week or more than 3 on an occasion for women.

The biggest predictor of transitioning to at-risk alcohol use was younger age, particularly among participants who were under the [drinking](#) age at the time of the first survey. Other factors were being male, not married, becoming divorced or separated, being in the military, being in good or excellent [health](#), smoking, drug use, and having an alcohol use disorder. The researchers found predictors of not transitioning to at-risk use were being black, reporting more stressful life experiences, having children

between the first and second rounds of the survey, and unemployment.

Predictors of continuing to drink too much were also being younger, male, having an [alcohol](#) use disorder, and using tobacco or other drugs. Being Black and/or Hispanic, receiving [alcohol use disorder](#) treatment, and having children between the two rounds of the survey were predictors of transitioning to lower-risk use.

"These findings suggest that not only do many people who drink, drink amounts associated with [health consequences](#), but that without intervention they are likely to continue to do so," Saitz says. "Screening or self-assessments, and counseling, feedback, or [public health](#) messaging have roles in interrupting these patterns. The predictors we identified may help target those efforts."

Provided by Boston University School of Medicine

Citation: Almost half of US adults who drink, drink too much, and continue to do so (2018, July 17) retrieved 16 April 2024 from <https://medicalxpress.com/news/2018-07-adults.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--