

Majority of those in residential care have advance directives

July 25 2018



(HealthDay)—More than three-quarters of those living in residential

care facilities have an advance directive, according to a *QuickStats* report published in the July 20 issue of the U.S. Centers for Disease Control and Prevention's *Morbidity and Mortality Weekly Report*.

Jessica Penn Lendon, Ph.D., from the National Center for Health Statistics in Hyattsville, Md., and colleagues used data from the National Study of Long-Term Care Providers (2016) to assess the percentage of residential care community residents with an advance directive and examine geographic differences.

The researchers found that overall, 77.9 percent of residents in [residential care](#) communities had an advance directive documented in their files. Geographically, by Census division, the highest percentage of residents who had an advance directive were located in the Mountain division (87.8 percent), followed by residents in East North Central (83.7 percent), New England (80 percent), West North Central (78.9 percent), Pacific (77.6 percent), South Atlantic (77.4 percent), East South Central (76.4 percent), Middle Atlantic (68.8 percent), and West South Central (64.9 percent).

More information: [Abstract/Full Text](#)

Copyright © 2018 [HealthDay](#). All rights reserved.

Citation: Majority of those in residential care have advance directives (2018, July 25) retrieved 20 March 2024 from <https://medicalxpress.com/news/2018-07-majority-residential-advance.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--