


Toward a better understanding of Parkinson's disease

July 4 2018



Immunohistochemistry for alpha-synuclein showing positive staining (brown) of an intraneural Lewy-body in the Substantia nigra in Parkinson's disease. Credit: Wikipedia

A new study, published today in Nature Structural and Molecular



Biology, moves researchers closer to understanding one of the crucial proteins involved in Parkinson's disease.

Parkinson's disease is the second most common neurodegenerative disease and commonly affects patients over 60 years of age. It is caused by the degeneration of neurons that are located in the mid-brain and are involved in the dopamine system. One of the causes of the disease is the failure to clear away the damaged mitochondria within these neurons. Mitochondria are the powerhouse of the cells, creating energy for the cell.

Mutations in a protein, parkin, are responsible for a genetic form of the disease. Now, by using powerful beams of x-rays, a McGill-based research team, led by Kalle Gehring, has been able to visualize the steps of activation of this crucial protein. With funding from the Michael J Fox Foundation and the Canadian Institutes of Health Research, the team is learning how parkin is regulated within cells. This research offers hope for using <u>parkin</u> as a target for <u>therapeutic agents</u> to slow or prevent the progression of Parkinson's disease.

More information: Véronique Sauvé et al. Mechanism of parkin activation by phosphorylation, *Nature Structural & Molecular Biology* (2018). DOI: 10.1038/s41594-018-0088-7

Provided by McGill University

Citation: Toward a better understanding of Parkinson's disease (2018, July 4) retrieved 1 May 2024 from https://medicalxpress.com/news/2018-07-parkinson-disease.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is



provided for information purposes only.