

What you eat while pregnant may affect your baby's gut

July 4 2018



Credit: CC0 Public Domain

A mother's diet during pregnancy may have an effect on the composition of her baby's gut microbiome—the community of bacteria living in the gut—and the effect may vary by delivery mode, according to study

published in the open access journal *Microbiome*.

Sara Lundgren, lead author of the study said: "Our study demonstrates an association of a readily modifiable factor, [maternal diet](#), with the infant gut microbiome. This knowledge may be key for developing evidence-based dietary recommendations for pregnant and lactating women."

Lundgren and colleagues at Dartmouth-Hitchcock medical Center, USA found that the gut microbiome in infants six weeks after [delivery](#) was composed mostly of Enterobacteriaceae (~20%), *Bifidobacterium* (18.6%), *Bacteroides* (10.44%) and *Streptococcus* (8.10%).

The authors identified three distinct clusters of microbes in the guts of the 97 babies included in this study that had been vaginally delivered. Cluster 1 was characterized by a high abundance of *Bifidobacterium*, cluster 2 showed a high abundance of *Streptococcus* and *Clostridium*, while cluster 3 had a high abundance of *Bacteroides*. These clusters were different for the 48 babies delivered by caesarian section, where cluster 1 showed a high abundance of *Bifidobacterium*, cluster 2 was characterized by high *Clostridium* but low *Streptococcus* abundance, and cluster 3 showed a higher abundance of Enterobacteriaceae.

The authors also observed what appear to be effects of certain aspects of the mothers' diets on the babies' gut microbiome. In babies delivered vaginally, the odds of being in cluster 2 were 2.73 times higher for each additional serving of fruit consumed by the mothers per day.

Bifidobacterium was found to be decreased in vaginally born infants if mothers ate more fruit, but increased in babies born by caesarian in relation to the mothers' consumption of red and processed meat. In babies delivered by caesarian section, the odds of being in [cluster](#) 2 were 2.36 higher for each additional maternal serving of dairy per day.

Lundgren said: "We analyzed infants delivered vaginally and by cesarean

section in separate groups due to our previous knowledge of the transfer of maternal microbiota to the infant that occurs during vaginal delivery, but not with [cesarean section delivery](#). We expected results to differ based on delivery mode, but we were surprised to find that the abundances of some microbes were increased in association with maternal intake of a food group in one delivery mode group, but decreased in the other delivery mode group."

To shed further light on the mechanisms by which maternal diet may affects children's health via the gut [microbiome](#) the authors used stool samples from 145 infants enrolled in the New Hampshire Birth Cohort Study, a research project that investigates how various factors affect the health of pregnant women and their children. Most infants whose data was used in this study were born vaginally (66.9%) and exclusively breast fed (70.3%) at six weeks of age. Information on the mothers' diets during pregnancy at 24 to 28 weeks gestation was obtained via food frequency questionnaires.

The authors caution that as their sample only included mothers and [infants](#) from Northern New England which has a relatively homogenous population, the generalizability of the findings may be limited. The authors also point out that the effects observed in this study may be due in part to maternal diet during lactation. The observational nature of the study does not allow for conclusions about cause and effect or the directionality of the observed association between maternal [diet](#) and the baby [gut microbiome](#).

More information: Sara N. Lundgren et al, Maternal diet during pregnancy is related with the infant stool microbiome in a delivery mode-dependent manner, *Microbiome* (2018). [DOI: 10.1186/s40168-018-0490-8](#)

Provided by BioMed Central

Citation: What you eat while pregnant may affect your baby's gut (2018, July 4) retrieved 25 April 2024 from <https://medicalxpress.com/news/2018-07-pregnant-affect-baby-gut.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.