

Preoperative opioids used by 23.1 percent of surgical patients

July 17 2018



(HealthDay)—Preoperative opioid use is reported in 23.1 percent of

patients undergoing surgery, according to a study published online July 11 in *JAMA Surgery*.

Paul E. Hilliard, M.D., from the University of Michigan Health System in Ann Arbor, and colleagues examined the prevalence of preoperative [opioid](#) use in a cross-sectional, observational study of [patients](#) undergoing surgery. Data were included for 34,186 patients.

The researchers found that preoperative opioid use was reported in 23.1 percent of patients. Hydrocodone bitartrate, tramadol hydrochloride, and oxycodone hydrochloride were the most common opioids used (59.4, 21.2, and 18.3 percent, respectively). Independent associations with preoperative opioid use were seen for ages 31 to 40 years (adjusted odds ratio [aOR], 1.26), tobacco use (former use aOR, 1.32; current use aOR, 1.62), illicit drug use (aOR, 1.74), higher pain severity (aOR, 1.33), depression (aOR, 1.22), higher Fibromyalgia Survey scores (aOR, 1.06), lower life satisfaction (aOR, 0.95), and more medical comorbidities (American Society of Anesthesiology score aOR, 1.47; Charlson Comorbidity Index aOR, 1.29). Patients undergoing orthopedic and neurosurgical spinal procedures most commonly reported preoperative opioid use (65.1 and 55.1 percent, respectively), while those undergoing thoracic procedures least commonly reported use (15.7 percent).

"These data provide important insights into this complicated patient population that would appear to help guide future preoperative optimization and perioperative opioid-weaning interventions," the authors write.

Several authors disclosed financial ties to the pharmaceutical industry.

More information: [Abstract/Full Text \(subscription or payment may be required\)](#)
[Editorial \(subscription or payment may be required\)](#)

Copyright © 2018 [HealthDay](#). All rights reserved.

Citation: Preoperative opioids used by 23.1 percent of surgical patients (2018, July 17) retrieved 27 April 2024 from

<https://medicalxpress.com/news/2018-07-preoperative-opioids-percent-surgical-patients.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.