

## Brief safety plan intervention in ER can cut suicidal behavior

July 17 2018

---



(HealthDay)—Use of the Safety Planning Intervention (SPI) plus follow-

up phone calls for suicidal patients presenting in the emergency department cuts suicidal behavior and increases the likelihood of outpatient mental health treatment over the next six months, according to a study published online July 11 in *JAMA Psychiatry*.

Barbara Stanley, Ph.D., from Columbia University in New York City, and colleagues assessed whether the SPI administered in nine Veterans Affairs emergency departments with telephone follow-up contact for suicidal [patients](#) was associated with reduced suicidal behavior and improved outpatient treatment engagement in the high-risk, six-month period following discharge.

The researchers found that the 1,186 patients in the SPI+ group were less likely to engage in [suicidal behavior](#) during the six-month follow-up period (3.03 percent) versus the 454 patients receiving usual care (5.29 percent). Suicidal behaviors were cut nearly in half with SPI+ (odds ratio, 0.56; P = 0.03). Additionally, patients receiving the SPI+ had more than double the odds of attending at least one outpatient mental health visit (odds ratio, 2.06; P

"The Safety Planning Intervention with structured follow-up telephone contact may be an effective brief suicide prevention [intervention](#) that can be implemented in [emergency](#) departments," the authors write.

**More information:** [Abstract/Full Text](#)

Copyright © 2018 [HealthDay](#). All rights reserved.

Citation: Brief safety plan intervention in ER can cut suicidal behavior (2018, July 17) retrieved 26 April 2024 from <https://medicalxpress.com/news/2018-07-safety-intervention-er-suicidal-behavior.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.