


# Spleen microbes of wild animals change with tick-borne illness

July 5 2018



Differentially abundant bacterial taxa in the spleen microbiota between

Anaplasma-positive and Anaplasma-negative samples identified by LEfSe analysis. Cladogram shows the highly represented taxa in Anaplasma-positive and Anaplasma-negative samples, respectively. A: Anaplasma-positive, in red; N\_A: Anaplasma-negative, in green. Credit: Ge, et al. 2018 / CC0

Anaplasmosis, a tick-borne febrile disease, can be carried by wild mammals before being transmitted to humans through a tick bite. Now, researchers reporting in *PLOS Neglected Tropical Diseases* have found that *Anaplasma* bacteria alter the patterns of other microbes in the spleens of mice and shrews.

The spleen, in mammals, acts as a blood filter. While studies have looked at the microbiota of the blood of mammals, few studies have been carried out on spleen microbiota. Additionally, it was unknown whether there were any vector-borne bacteria in Chongming Island, the third biggest island in China and where the study was conducted.

Yan Ge, of Tongji University, China, and colleagues trapped 35 mice and shrews on Chongming Island. They then sequenced the DNA isolated from spleen samples of each animal. The genes allowed them to detect [bacterial phyla](#) and hierarchies in each animal's spleen.

The most bacteria detected in the animals' spleens included *Anaplasma*, *Rickettsia*, and *Coxiella*, all [tick-borne diseases](#) that can infect humans. Interestingly, animals that were Anaplasma-positive had a significantly different overall bacterial composition in their [spleen](#) compared to animals that were Anaplasma-negative. Within Anaplasma-positive [animals](#), there were two smaller clusters with similar bacterial features, likely linked to a separate independent factor not analyzed in this study. More subtle patterns in microbiota also emerged based on animal gender, type, and geographic location.

"The detection of vector-borne bacteria highlights the role of wild mice and shrews as animal reservoirs with potential public health importance in the studied areas," the researchers say.

**More information:** *PLOS Neglected Tropical Diseases* (2018). [DOI: 10.1371/journal.pntd.0006499](https://doi.org/10.1371/journal.pntd.0006499)

Provided by Public Library of Science

Citation: Spleen microbes of wild animals change with tick-borne illness (2018, July 5) retrieved 6 May 2024 from <https://medicalxpress.com/news/2018-07-spleen-microbes-wild-animals-tick-borne.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--