

Can community exercise prevent bone loss from weight loss in older adults?

August 8 2018



Credit: CC0 Public Domain

In a *Journal of Bone and Mineral Research* study of older adults with obesity who were cutting calories, an intervention that incorporated resistance training, aerobic training, or neither did not prevent bone loss

associated with active weight loss. The study's results suggested that resistance training may help minimize long-term hip bone loss, however.

"If minimizing [bone loss](#) during active weight loss proves necessary to offset long-term skeletal fragility, then our results suggest that resistance exercise may need to be coupled with other intervention strategies to maximize skeletal benefit," said lead author Dr. Kristen Beavers, of Wake Forest University.

Additional research should seek to elucidate the mechanisms underlying weight-loss-induced bone loss, so that safe and effective strategies can be designed to preserve all aspects of bone health in dieting older individuals.

More information: Kristen M Beavers et al, Effect of Exercise Modality During Weight Loss on Bone Health in Older Adults With Obesity and Cardiovascular Disease or Metabolic Syndrome: A Randomized Controlled Trial, *Journal of Bone and Mineral Research* (2018). [DOI: 10.1002/jbmr.3555](https://doi.org/10.1002/jbmr.3555)

Provided by Wiley

Citation: Can community exercise prevent bone loss from weight loss in older adults? (2018, August 8) retrieved 26 April 2024 from <https://medicalxpress.com/news/2018-08-bone-loss-weight-older-adults.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.