


Brain-heart activity predicts post-malaria epilepsy in mice

August 27 2018



Credit: Bahari et al., *JNeurosci* (2018)

Animals that develop epilepsy after an infection can be identified as early as three months prior to their first seizure by measuring interactions between the brain and the heart, according to new research using a mouse model of post-cerebral malaria epilepsy. Published in *JNeurosci*, this finding could inform efforts to diagnose and treat acquired epilepsy.

Cerebral malaria afflicts more than three million people worldwide, affects [young children](#), and leads to epilepsy in an estimated 15 percent

of survivors. Reducing the risk of developing epilepsy and associated fatal complications, such as sudden unexplained death in epilepsy, requires a reliable way to detect and monitor epileptogenesis.

By recording neural and cardiac activity in a [mouse model](#) of cerebral malaria, Fatemeh Bahari, Bruce Gluckman and colleagues discovered a signal transmitted between the brain and the heart that occurred only in the 75 percent of mice that acquired epilepsy. Translating this biomarker to humans has the potential to improve therapeutic approaches in patients at risk of developing epilepsy, such as those recovering from [traumatic brain injury](#) or stroke.

More information: *JNeurosci* (2018). [DOI: 10.1523/JNEUROSCI.1130-18.2018](#)

Provided by Society for Neuroscience

Citation: Brain-heart activity predicts post-malaria epilepsy in mice (2018, August 27) retrieved 25 April 2024 from <https://medicalxpress.com/news/2018-08-brain-heart-post-malaria-epilepsy-mice.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
